

V O Y A G E

OF

H. M. S. BLONDE

TO THE

SANDWICH ISLANDS,

IN THE YEARS 1824-1825.

CAPTAIN THE RIGHT HON. LORD BYRON, COMMANDER.

LONDON:

JOHN MURRAY, ALBEMARLE-STREET.

MDCCKXXVI.

6. ; *native name*, Erepeio.

Muscicapa Sandwichensis. Linn.

Sandwich Flycatcher.

7. *Fringilla Rufa*; *native name*, Akepakepa.

Length four inches and a quarter; bill hard, straight, short, and conical; three-eighths of an inch in length, sharp-pointed; body rufous; tail and wings brownish; toes and legs strong, formed for perching; black coloured tongue, short and tubular, divided into filaments at the end.

8. *Fringilla Sandwichensis*; *native name*,

A. Length, five inches; whole of the back dull olive green; greater and lesser wing coverts tipped with dirty white; wings and tail brown, edged with green; belly greenish white; bill straight, sharp-pointed, half an inch long; tongue bifid.

B. Differing, in being of a much lighter colour.

9. ; *native name*, Amauii.

Turdus Sandwichensis. Linn.

Sandwich thrush.

Found chiefly in Hawaii. There is a variety of the same at Oahu.

10. *Strix Sandwichensis*; *native name*, Puaho.

Length thirteen inches. Mottled all over with dirty white and reddish brown.

11. *Corvus Tropicus*. Linnæus.

12. ; *native name*, Alai.

Fulica Chloropus. Linn.

Common Moor-hen.