

S U P P L E M E N T U M
I N D I C I S O R N I T H O L O G I C I ,

S I V E

S Y S T E M A T I S O R N I T H O L O G I Æ .

S T U D I O E T O P E R A

J O A N N I S L A T H A M , M . D . S . R . S .

L O N D I N I :

P R O S T A T A P U D G . L E I G H , J . E T S . S O T H E B Y ,
Y O R K - S T R E E T , C O V E N T - G A R D E N .

M , D C C C , I .

38.
F.

Sonninen-F. cauda subforcipata, corpore cinereo subtus albo, tibiis plumosis.
fis. *Sonnini.*

Sonnini's Falcon, *Gen. Syn. Sup.* ii. p. 52. No. 59.

HABITAT in *Agypto* satis frequens: longit. fere ped. 1. :—*rostrum*,
cera, *irides*, & *pedes* flavi: *rectrices* intermediae paulo breviores: *remiges*
ultra apicem caudae extendunt: *sinuiput* album.

39.
F.

melanops. F. supra nigricans, subtus ferrugineus lineis cinereis transversis, orbitis
nigris.

Black-eyed Falcon, *Gen. Syn. Sup.* ii. p. 52. No. 60.

HABITAT in *Nova Hollandia*: long. poll. 34. :—*caput* & *collum* cinerea:
circa oculos rufum, *orbitis* nigris: *rostrum* nigrum.

40.
F.

radiatus. F. nigro radiato-maculatus, corpore ferrugineo, alis caudaque elongata
fuscis.

Radiated Falcon, *Gen. Syn. Sup.* ii. p. 53. No. 61. tab. 121.

HABITAT in *Nova Hollandia*: longit. poll. 22. :—*corpus* totum laete
ferrugineum, fasciis maculisque difformibus nigris: *cauda* longissima ro-
tundata, fasciis 8, 9 nigris: *rostrum* nigrum: *pedes* caerulei.

41.
F.

connivens. F. badius subtus flavescens maculis fuscis, cervice & axillaribus albo
maculatis.

Winking Falcon, *Gen. Syn. Sup.* ii. p. 53. No. 62.

HABITAT in *Nova Hollandia*, magnitudine F. *Cyanei*: *remiges* &
rectrices badio & albo fasciatæ: *tibiae* plumis tectæ: *rostrum* pallidum:
irides et *digiti* flavi.

42.
F.

discolor. F. griseo-nigricans subtus ferrugineus, rectricibus inferioribus alarum
& caudae albis.

Falco discolor, *Aët. Soc. N. H. de Paris*, i. pl. 1. p. 911. 4.


Der bunt Falke, *Allg. U. d. Vog.* ii. S. 680. 128.

Rusty and grey Falcon, *Gen. Syn. Sup.* ii. p. 53. No. 63.

HABITAT in *Cayana*.

F. fuscus

SUPPLEMENT. II.
to the
GENERAL SYNOPSIS
of
BIRDS.



L O N D O N.
Printed for Leigh, Sotheby & Son,
York Street, Covent Garden.

M D C C C I.

THIS species is twenty-two inches long, and four feet broad from the tip of one wing to that of the other: the bill is black: the cere and space round the eye pale blue: irides brown: the plumage in general of a fine rust colour, spotted and streaked all over with black, but crossed on the head and neck with fine lines of the same: wings and tail brown, marked with blackish or dusky bars: the tail is very long, with eight or nine black bars, the wings reaching to the middle of it: the legs are blue; claws long, black, and sharp.

This inhabits *New Holland*, but is probably a scarce species, only one having been met with, which was found nailed to the side of a barn.

61.
RADIATED
F.
DESCRIPTION.

PLACE.

62.
WINKING
F.
DESCRIPTION.

SIZE of the *Ringtail*; length eighteen or nineteen inches: bill pale, with a black point: irides yellow: the general colour of the plumage is a deep chocolate brown, spotted with rusty white on the lower part of the neck behind, and on the axillaries of the wings: the quills are obliquely, and the tail feathers transversely barred with the same: the under parts as far as the breast, dirty yellowish white, with short dusky streaks: legs covered to the toes with pale ash-coloured feathers.

Inhabits *New Holland*, but no history annexed, further than that it has a wonderful faculty of contracting and dilating the iris; and that the native name is *Geora-a-Gang*.

PLACE.

Falco discolor, *AH. Soc. N. H. de Paris*, i. pl. 1. p. 911. 4.
Der bunte Falke, *Allg. U. d. Vog.* ii. S. 683. 128.

63.
RUSTY and
GREY
F.
DESCRIPTION.

THE size of this bird is not mentioned: but the description observes, that the upper parts of the plumage are dusky grey; the under, breast, belly, and thighs, dusky rust colour: the under wing coverts and those of the tail white.

Inhabits *Cayenne*.

PLACE.