

3. *NINOX MINDORENSIS*, sp. n.

An adult female of this little Mindoro Hawk-Owl was obtained in the lowlands, and is nearly allied to *N. spilocephala*, Tweedd., from Mindanao and Basilan, resembling that species in having the top of the head and nape barred with buff. It may, however, be at once recognized by having the whole of the underparts, including the belly and flanks, tawny buff, transversely barred with brown, while in the female of *N. spilocephala*, though the breast is generally like that of the present species, the belly and flanks are always white, with longitudinal reddish-brown shaft-stripes. Total length 8·0 inches, wing 6·0, tail 3·0, tarsus 1·1.

I have of course carefully compared this species with Messrs. Bourns and Worcester's description of *N. spilonotus*, specimens of which were obtained in Mindoro, but the much smaller size of this bird serves to at once distinguish it.

4. *NINOX JAPONICA* (Temm. & Schl.) ; Grant, Ibis, 1896, p. 111.

A male example of the Japanese Hawk-Owl was procured. It is a nearly typical example of *N. japonica*, much darker than the bird from Fuga Island, described in the last paper, the predominating colour of the underparts being reddish brown.

5. *CORVUS PUSILLUS*, Tweeddale, P. Z. S. 1878, p. 622.

Mr. Whitehead sends several examples of this small Philippine Raven from the lowlands of Mindoro. They are all in splendid freshly moulted plumage, the wings especially having a fine deep purplish gloss.

6. *ORIOLOUS CHINENSIS*, Linn. ; Grant, Ibis, 1896, p. 111.

A female of the Chinese Oriole from the lowlands.

7. *ARTAMIDES MINDORENSIS*, Steere, List of Birds & Mamm. Philippines, p. 14 (1890).

Both sexes of the Mindoro Cuckoo-Shrike are represented.

8. *MUSCICAPULA LUZONIENSIS*, Grant, Ibis, 1894, p. 505.

Mindoro examples of the Luzon Flycatcher are perfectly similar to those obtained in the highlands of Benguet and