

Table (continued).

	Salvadori, <i>op. cit.</i>	Locality.	No. of examples.
VI. GALLINÆ.			
42. <i>Megapodius tenimberensis</i> , sp. nov.		Loetoe.	2
VII. GRALLATORES.			
43. <i>Orthorhampus magnirostris</i>	iii. p. 290	Kirimoen.	1
44. <i>Charadrius fulvus</i>	iii. p. 294	Maroe.	1
45. <i>Ægialitis geoffroii</i>	iii. p. 298	Maroe.	1
46. <i>Lobivanellus miles</i>	iii. p. 306	Larat.	1
47. <i>Totanus incanus</i>	iii. p. 320	Moloe.	1
48. <i>Numenius variegatus</i>	iii. p. 332	Larat.	1
49. <i>Ardea sumatrana</i>	iii. p. 340	Larat.	1
50. <i>Demigretta sacra</i>	iii. p. 345	Larat.	1
VIII. NATATORES.			
51. <i>Nettapus pulchellus</i>	iii. p. 385	Larat.	1
52. <i>Dendrocygna guttata</i>	iii. p. 388	Larat.	1
53. <i>Tadorna radjah</i>	iii. p. 391	(No ticket.)	1
54. <i>Onychoprion anæsthetus</i>	iii. p. 449	Moloe.	1

I will now give descriptions of the new species, and notes upon several others imperfectly known.

4. *NINOX FORBESI*, sp. nov. (Plate XI.)

Supra rufescenti-brunnea, fere unicolor, in alarum tectricibus et scapularibus fasciis albis variegata; fronte et superciliis albis; alarum remigibus terreno-brunneis, nigro transfasciatis; subtus dorso concolor, mento albicante, ventre albo transfasciato; tarsi, omnino plumosis, cum subalaribus rufis unicoloribus; alarum et caudæ pagina inferiore pallide corylino-brunnea nigro regulariter transfasciata; rostri nigri apice flavicante; digitis fuscis setis obtectis: long. tota 11·0, alæ 7·4, caudæ 4·5, tarsi 1·3.

Hab. Loetoe, Timor Laut.

Obs. Sp. quoad colores *N. hantu* maxime affinis, sed facie alba fasciis ventris albis, et alis subtus nigro vittatis diversa.

The single specimen of this Owl is a male, obtained at Loetoe on August 9, 1881. It is noted:—"Irides golden; bill pale cinereous; feet pale yellow, covered with bristly hairs; soles of feet nearly orange."

I have dedicated this apparently distinct species to its discoverer, Mr. Henry Ogg Forbes, F.Z.S.

5. *STRIX SORORCULA*, sp. nov.

Supra terreno-fusca flavicante variegata, et punctis rotundis albis regulariter aspersa; disco faciali amplo albo, margine nigricanti-brunneo circumdato; macula anteoculari nigricante; remi-