under surface of wings and tail rufous white, spotted or banded with black.

Length 16", wings 8" 9".

East Gilolo.

Probably a very young stage of Astur griseogularis.

*Accipiter erythrauchen.

Slaty-black; lore, cheeks, mentum, abdomen, and under tail-coverts slaty-white; side of neck and nape castaneous rufous; breast, sides of abdomen, and thighs whitish-rufous; under wing-coverts rufous-white, varied with slaty-white; quills beneath banded with rufous or slaty-white and slaty-black.

Length 11" 9", wings 8" 9". East Gilolo (Wall. Coll.).

MICRONISUS SOLOENSIS.

Falco soloensis, Horsf. Linn. Trans. xiii. p. 137.

Falco cuculoides, Temm. Pl. Col. 110. 129.

Tachyspiza soloensis, Kaup. Classif. der Saug. und Vög. p. 116. Micronisus soloensis, G. R. Gray.

Batchian (Wall. Coll.).

STRIGIDÆ.

*ATHENE RUFOSTRIGATA.

This bird approaches the Athene connivers in general appearance, but it is of a blackish-slate colour on the upper surface; the bands on the quills and tail-feathers are less prominent, and the white spots on the wings are less numerous; while the longitudinal streaks on the white under surfaces are of a rufous colour, slightly varied with slaty-black.

Length 17" 9", wings 11" 6". East Gilolo (Wall. Coll.).

ATHENE HYPOGRAMMA.

B.M.

Upper surface uniform deep rufous-brown, more obscure on the head; scapulars and some of the wing-coverts banded with white; front and throat white; under surface entirely white, broadly banded with brownish-rufous; tail banded with paler colour; tarsi plumed and rufous-white; toes covered only with strong scattered hairs.

Length 15'' 3''', wings 8'' 9'''.

Batchian and Gilolo.

This bird bears a great similarity to the Athene variegata (Q. & G.), but it does not exhibit the marks visible on the upper surface of that bird and it also differs in having the barring of the under surface extending on the tail-coverts. It is also larger in all its proportions.

Athene squamipila, Pr. B. Ceram.

EPHIALTES LEUCOSPILA.

B.M.

Upper surface rufous, speckled and irregularly striated longitu-

dinally with black; the marks on the head are broader and more defined; under surface varied with rufous and white, with black irregular marks down the shaft of each feather; scapulars, wing-coverts, outer margins of quills, and of the outer tail-feather spotted with white; tarsi plumed; toes entirely naked.

Length 11", wings 6" 6".
Batchian and Eastern Gilolo.

In general appearance this bird is very like *Ephialtes manadensis* (Q. & G.), but it is larger in all its proportions, and is more prominently marked with white on the wing-coverts.

Ephialtes magicus, Müll. Amboyna; Banda Islands.

CAPRIMULGIDÆ.

BATRACHOSTOMUS PSILOPTERUS.

Rufous-cinnamon; front, middle of throat, breast, abdomen, and under tail-coverts varied with rufous and white—the latter colour margined with black; wing-coverts, scapulars, and some of the tertials spotted with white, surrounded with black; a spot behind each eye and at the base of mandibles white; tail irregularly banded with black; quills black, with the outer web cinnamon colour; bristles lengthened and deep black.

Length 12", wings 7".

Batchian and Gilolo (Wall. Coll.).

HIRUNDINIDÆ.

MACROPTERYX MYSTACEUS.

B.M.

Cypselus mystaceus, Less. Voy. de la Coqu. Zool. i. t. 22. Macropteryx mystaceus, Swains. Classif. of B. ii. p. 340. Dendrochelidon mystaceus, Boié, Isis, 1844, p. 166. Batchian and Eastern Gilolo; Amboina.

COLLOCALIA HYPOLEUCA.

B.M.

Collocalia hypoleuca, G. R. Gray, Proc. Z. S. 1858, p. 170. Batchian.

Collocalia esculenta (?). Amboyna.

HIRUNDO GUTTURALIS.

B.M.

Hirundo gutturalis, Scop. Hirundo panayensis, Gmel. S. N. i. p. 1018. Hirundo javanica, Sparrm. Mus. Carls. t. 100. Batchian and Gilolo.

CORACIADÆ.

EURYSTOMUS ORIENTALIS.

Coracias orientalis, Linn. S. N. i. p. 159. Eurystomus orientalis, Steph. Gen. Zool. xiii. p. 99. Colaris orientalis, Cuv. Règ. Anim. 1817, i. p. 401. Batchian and Ternate (Wall. Coll.).