

	LOMBOCK.	FLORES.	TIMOR.
159. <i>Megapodius reinwardti</i> , Wagl.....	<i>M. reinwardti</i>	<i>M. reinwardti</i>
160. <i>Charadrius longipes</i> , Temm.....	<i>C. longipes</i>	<i>C. longipes</i> .
161. — <i>leschenaultii</i> , Pall.....	<i>C. leschenaultii</i>
162. <i>Totalanus hypoleucus</i> , Temm.....	<i>T. hypoleucus</i>	<i>T. hypoleucus</i> .
163. — <i>glareola</i> , Gm.....	<i>T. glareola</i> .
164. <i>Numenius uropygialis</i> , Gould.....	<i>N. uropygialis</i>
165. <i>Scolopax horsfieldi</i> , Gray.....	<i>S. horsfieldi</i> .
166. <i>Himantopus leucocephalus</i> , Gould.....	<i>H. leucocephalus</i> .
167. <i>Ardea typhon</i> , Temm.....	<i>A. typhon</i>
168. <i>Egretta immaculata</i> , Gould.....	<i>E. immaculata</i>
169. — <i>syrmatophora</i> , Gould.....	<i>E. syrmatophora</i> .
170. — <i>nigrirostris</i> , Gray.....	<i>E. nigrirostris</i> .
171. <i>Herodias coromanda</i> , Bodd.....	<i>H. coromanda</i> .
172. — <i>jugularis</i> , Blyth.....	<i>H. jugularis</i> .
173. — <i>novæ-hollandiæ</i> , Lath.....	<i>H. novæ-hollandiæ</i> .
174. <i>Butorides javanica</i> , Horsf.....	<i>B. javanica</i>	<i>B. javanica</i> .
175. <i>Ardeola sinensis</i> , Gm.....	<i>A. sinensis</i>
176. <i>Rallina phœnicura</i> , Temm.....	<i>R. phœnicura</i> .
177. — <i>philippensis</i> , L.....	<i>R. philippensis</i> .
178. <i>Porzana quadristrigata</i> , Horsf.....	<i>P. quadristrigata</i> .
179. <i>Gallinula orientalis</i> , Horsf.....	<i>G. orientalis</i>
180. — <i>frontata</i> , Wall.....	<i>G. frontata</i>
181. <i>Porphyrio smaragdinus</i> , Temm.....	<i>P. smaragdinus</i> .
182. <i>Anas gibbifrons</i> , Müll.....	<i>A. gibbifrons</i>	<i>A. gibbifrons</i> .
183. — <i>superciliosa</i> , Gould.....	<i>A. superciliosa</i> .
184. <i>Dendrocygna vagans</i> , Eytton.....	<i>D. vagans</i> .
185. <i>Podiceps tricolor</i> , G. R. G.....	<i>P. tricolor</i>	<i>P. tricolor</i> .
186. <i>Graculus melanoleucus</i> , Vieill.....	<i>G. melanoleucus</i> .
	63 species.	86 species.	118 species.

ACCIPITER SYLVESTRIS.

Cinereus, capite pallidiore, gula pallide rufescente, pectore et abdomine rufis anguste albo fasciatis, alis subtus et tectricibus caudæ inferioribus albescentibus, rectricibus ad basin albis plus minusve nigro fasciatis.

Ashy above, rufous beneath, narrowly banded with whitish ; wings white on the under surface, except the ends of the quills ; tail white at the base, and the feathers towards the centre with eight or nine blackish bands. This bird resembles *A. iugularis* in wanting the rufous collar, but is much smaller and paler-coloured ; it is also like *A. cruentus* in its banded under surface, but, though smaller, has a larger bill than that species. Bill black ; cere and legs yellow.

♂. Total length $12\frac{1}{2}$ inches ; wings 7 inches ; tail $5\frac{1}{2}$ inches.

♀. Total length $13\frac{3}{4}$ inches ; wing 8 inches ; tail $6\frac{1}{4}$ inches.

Hab. Flores.

SCOPS SILVICOLA.

Rufo-fuscus, plumis nigro lineatis et maculatis ; tectricibus alarum albo marginatis ; remigibus fuscis, fasciis plurimis externe pallide rufis, interne obsoletis ; cauda obscura, rufo irrorata, fasciis decem rufis ; tarsis dense vestitis, digitis nudis.

Pale dusky rufous, the feathers irrorated with dusky and with a median dark line; greater and middle wing-coverts white-tipped; quills dusky, with obsolete paler bands, on the outer web with pale rufous bands; tail with about ten narrow rufous bands, which are dusky-margined; facial setæ very long; tarsi densely feathered; toes naked.

Total length 12 inches; wing $8\frac{1}{2}$ inches; tail $4\frac{1}{2}$ inches; bill, from gape, $1\frac{1}{10}$ inch.

Hab. Flores.

Remark.—The only specimen obtained was immature; but it seems very distinct from all other Malayan species.

ATHENE FLORENSIS.

Fusco-brunnea, subtus alba, brunneo maculato-striata; fronte gulaque albescensibus; cauda fasciis quinque obscurioribus, rectricibus albo terminatis.

Darker and much larger than the other Malayan forms of *Athene hirsuta*.

Total length $12\frac{3}{4}$ inches; wing $9\frac{1}{4}$ inches; tail $5\frac{1}{4}$ inches; tarsus and mid toe, without claw, $2\frac{1}{4}$ inches; bill, from gape, 1 inch.

Hab. Flores.

LORICULUS FLOSCULUS.

Viridis, dorso obscuriore; macula gulari, uropyglio tectricibusque caudæ superioribus coccineis; macula nuchali fulvescente; alis caudaque subtus glauco-viridibus, rectricibus flavo terminatis.

Green, above darker; middle of the body beneath with a yellowish tinge; nape fulvous orange; elongate spot on throat red, rump and upper tail-coverts crimson; tail above dark green tipped with yellow and red; quills and tail-feathers beneath margined with verditer-blue; bill red; feet orange-red.

Total length $4\frac{3}{4}$ inches; wing 3 inches.

Hab. Flores.

Remarks.—This pretty little bird is allied to the *L. pusillus* of Java and *L. indicus* of Ceylon; but also approaches in coloration to *L. galgulus* and others of the black-billed group. A single specimen only was obtained.

ANTHUS MEDIUS.

A. australi similis, sed minor, ungue postico magis curvato: similis

A. malayano, sed rostro breviore, pectoris striis majoribus, ungue postico breviore et magis curvato.

Dusky or blackish above; the feathers margined with ochreish yellow or ashy, beneath white or yellowish; the breast and a line from each angle of the mouth dusky-spotted; a yellowish line above the eye; tail dusky, outer feather white, except the base, the next dusky on the inner margin; bill blackish horn, beneath yellowish; feet very pale; iris dark.