

Measurements.

No.	Locality	Sex	Wing	Tail	Tar- sus	Exposed culmen
836 ¹	Vic. Paramaribo	♂	196	270	44	46
837 ¹	" "	♂	195	253	43	46
838 ¹	" "	♂	195	260	42	43
1361 ¹	" "	♂	195	255	44	48
1362 ¹	" "	♂	199	265	43	48
839 ¹	" "	♀	191	240	43	46
1360 ¹	" "	♀	184	244	42	43

89. CROTOPHAGA ANI Linné.

Five specimens, both sexes, Vicinity of Paramaribo, and Lelydorp, April, May, and December.

BUBONIDAE.

90. PULSATRIX PERSPICILLATA PERSPICILLATA (Latham).

Six specimens, both sexes, Vicinity of Paramaribo, and Javaweg, January, March, April, and May.

The darker specimens are fully as dark above as *P. p. saturata* Ridgway, but the breast and belly are conspicuously paler buff and never show any cross barring.

Birds from Trinidad are decidedly paler, and we name the island form

90a. PULSATRIX PERSPICILLATA TRINITATIS, subsp. nov.

Type.—M. C. Z. 29,469. Adult. Trinidad. 1880? C. S. Cazabon.

Characters.—In general like *P. p. perspicillata* (Lath.), but upper parts and pectoral collar paler, not so sooty; breast and belly average slightly paler buff.

Measurements.—*Type*, wing, 305; tail, 165; culmen from cere, 28.

Remarks.—In color *P. p. trinitatis* is about intermediate between *P. p. perspicillata* (Lath.) and *P. p. sharpei* Berl., although its range is not geographically between the ranges of these two forms. The above confirms Hellmayr's views (Nov. zool., 1906, 13, p. 45).