

S U P P L E M E N T U M
I N D I C I S O R N I T H O L O G I C I ,

S I V E

S Y S T E M A T I S O R N I T H O L O G I Æ .

S T U D I O E T O P E R A

J O A N N I S L A T H A M , M . D . S . R . S .

L O N D I N I :

P R O S T A T A P U D G . L E I G H , J . E T S . S O T H E B Y ,
Y O R K - S T R E E T , C O V E N T - G A R D E N .

M , D C C C , I .

AVES ACCIPITRES. Strix.

27

HABITAT in *Africa*, magnitudine *Str. Otus*: corpus fuscum supra maculis rufo-nigricantibus & albis, subtus lunulis varium: facies albida, apicibus pennarum nigris.

6.

Str.

Huhula.

S. albo fasciata corpore cinereo, alis fuscis, cauda fasciis tribus albis.

Le Huhul, Levaill. Ois. i. p. 167. pl. 41.

Strix Huhula, Daudin. Orn. ii. p. 190.

Fasciated Owl, Gen. Syn. Sup. ii. p. 62. No. 11.

HABITAT in *Africa*, magnitudine *Str. Ulula*: corpus cinereum lineis numerosis transversis albis, subtus undulatum albido fasciatum; cauda cuneata fasciis tribus obliquis albis: rostrum, irides, & pedes flavi.

7.

Str.

griseata.

S. varia, corpore supra fusco rufoque, subtus rufo-alba, tectricibus alarum caudaque albo maculatis: superciliis albis.

La Chouette à Aigrette blanche, Levaill. Ois. i. p. 169. pl. 43.

Strix griseata, Daudin. Orn. ii. p. 207.

Supercilious Owl, Gen. Syn. Sup. ii. p. 63. No. 12.

HABITAT in *Guiana*, magnitudine *Str. Otus*: regio periopthalmica vix conspicua, supra oculos vitta alba ad axillas utrinque fere ducta: rostrum mediocre flavum: pedes nigricantes.

8.

Str.

Georgica.

S. fasciata capite lævi, corpore fusco subtus albo, regione periopthalmica albo annulata.

Georgian Owl, Gen. Syn. Sup. ii. p. 64. No. 13.

HABITAT in *Georgia Americana*: long. poll. 16½:—magnitudo *Str. Flammea*: rostrum flavum: gula & pectus fuscum fasciis albidis: abdomen albidum maculis fusco-rubris: femora albida punctis nigricantibus.

9.

Str.

Boobook.


S. capite lævi, corpore supra fusco subtus ferrugineo longitudinaliter maculato.

Boobook Owl, Gen. Syn. Sup. ii. p. 64. No. 14.

HABITAT in *Nova Hollandia*, magnitudine *Stridula*: corpus supra fusco-ferrugineum maculis flavescentibus: gula flava rufo maculata.

S. capite

SUPPLEMENT II.
to the
GENERAL SYNOPSIS
of
BIRDS.



L O N D O N.
Printed for Leigh, Sotheby & Son,
York Street, Covent Garden.

M D C C C I.

13.
GEORGIAN
O.

DESCRIPTION.

SIZE of the *Barn Owl*: length sixteen inches and a half: bill yellow: the plumage on the upper parts of the bird is brown, banded with yellowish: throat and breast pale brown, crossed with whitish bands: belly yellowish white, marked with longish red brown streaks: thighs and legs woolly, whitish or very pale in colour, varied with small blackish spots: quill and tail feathers brown, crossed with four or five white bands.

PLACE.

I met with this specimen in the collection of Mr. *Humbries*, said to have come from several miles within *South Georgia*, in *America*.

14.
BOOBOOK
O.

DESCRIPTION.

SIZE of the *Brown Owl*, and at first sight somewhat resembles it, especially on the upper parts; but marked with yellow streaks on the head, and with spots of the same on the back: the chin and throat are yellow, streaked and spotted with rufous: belly ferruginous, with pale irregular spots and markings: thighs and legs covered with yellow downy feathers, marked with black: toes of a brownish red: the bill is small and dusky.

PLACE.

This inhabits *New Holland*, where it is known by the name of *Boobook*.

15.
SPECTACLE
O.

VAR. A.

Str. perspicillata, *Ind. Orn.* i. p. 58. 24?

La Chouette à collier, *Lewinill. Ois.* i. p. 169. pl. 42.—*Daud. Orn.* ii. p. 193.

Spectacle Owl, *Var. Gen. Syn. Sup.* p. 5. pl. 107.

DESCRIPTION.

THE whole head in this bird is deep rufous, with a white throat, which colour passes between the eyes, and over them as an eyebrow; whereas in that figured in the *Gen. Synopsis*, *Sup.* pl. 107, the whole head is white, with a large rufous patch, in the middle of which