

392. **Tyto novaehollandiae novaehollandiae.**

Strix? novaehollandiae Stephens, in Shaw's *Gen. Zool.* vol. xiii. pt. ii. p. 61. 1826 : New South Wales.

Synonyms :

Strix personata Vigors, *Proc. Zool. Soc. (Lond.)* 1831. p. 60 : New South Wales.

Strix cyclops Gould, *Proc. Zool. Soc. (Lond.)* 1836. p. 140 (1837) : New South Wales.

Chestnut-faced Owl.

Mathews, *Handlist* No. 295.

Range : Queensland, New South Wales, Victoria, South Australia.

393. **Tyto novaehollandiae castanops.**

Strix castanops Gould, *Proc. Zool. Soc. (Lond.)* 1836. p. 140 (1837) : Tasmania.

Masked Owl.

Mathews, *Handlist* No. 296.

Range : Tasmania.

394. **Tyto novaehollandiae perplexa** subsp. n.

Western Chestnut-faced Owl.

Mathews, *Handlist* No. 295 (pars).

Differs from *T. n. novaehollandiae* in its larger size and darker coloration.

Type : West Australia (East Beverley), No. 4484.

Range : West Australia.

395. **Tyto novaehollandiae kimberli** subsp. n.

Pallid Chestnut-faced Owl.

Mathews, *Handlist* No. 295 (pars).

Differs from *T. n. novaehollandiae* in its much paler coloration and smaller size.

Type : East Kimberley, West Australia, No. 4485.

Range : North-West Australia.

396. **Tyto tenebricosa tenebricosa.**

Strix tenebricosus Gould, *Proc. Zool. Soc. (Lond.)* 1845. p. 80 : New South Wales (Clarence River).

Sooty Owl.

Mathews, *Handlist* No. 297 (pars).

Range : New South Wales.

397. **Tyto tenebricosa multipunctata** subsp. n.

Northern Sooty Owl.

Mathews, *Handlist* No. 297 (pars).

Differs from *T. t. tenebricosa* in its smaller size and more spotting above and below. Wing 263 mm. ; typ. w. 290 mm.

Type : Johnston River, North Queensland, No. 4276.

Range : North Queensland.