

**EMERGENCY PETITION TO LIST THE  
DELTA SMELT (*Hypomesus transpacificus*)  
AS AN ENDANGERED SPECIES  
UNDER THE ENDANGERED SPECIES ACT**



U.S. Fish and Wildlife Service photo

**Submitted To:** U. S. Fish and Wildlife Service, Washington D.C. and Sacramento Field Office, California

**Submitted By:** Center for Biological Diversity  
The Bay Institute  
Natural Resources Defense Council

**Date:** March 8, 2006

## EXECUTIVE SUMMARY

The Center for Biological Diversity, the Bay Institute and the Natural Resources Defense Council formally request that the U.S. Fish and Wildlife Service (USFWS) change the listing of the delta smelt (*Hypomesus transpacificus*) from threatened to endangered under the federal Endangered Species Act (ESA), on an emergency basis. In the past few years, the delta smelt population has plummeted to the lowest levels ever recorded and, based on population viability analysis, the species is in imminent danger of extinction, the criterion for endangered status. Recent surveys indicate that environmental conditions in the delta smelt's critical habitat, the upper San Francisco Bay/Sacramento-San Joaquin Delta Estuary, have also declined, threatening the planktonic food web upon which the species depends. The magnitude and frequency of occurrence of threats to the species identified by USFWS are also increasing. The petitioners are conservation organizations with an interest in protecting the delta smelt and its Delta habitat.

Delta smelt are endemic to the upper reaches of the Bay-Delta Estuary. The species requires specific environmental conditions (freshwater flow, water temperature, salinity) and habitat types (shallow open waters) within the estuary for migration, spawning, egg incubation, rearing, and larval and juvenile transport from spawning to rearing habitats. Delta smelt feed exclusively on plankton and most individual fish live only one year. Although they are restricted to a relatively small geographic range, delta smelt use different parts of the estuary at different life history stages. Throughout most of their life span, delta smelt inhabit low salinity habitat, at the interface of inflowing fresh water from the Sacramento and San Joaquin Rivers and salt water from the Pacific Ocean. Spawning adults, larvae, and young juveniles are found further upstream in the Delta, where they are vulnerable to lethal entrainment into federal, state, industrial, and local agricultural water diversion and export facilities.

As recently as thirty years ago, the delta smelt was one of the most common and abundant of the pelagic fishes in the estuary. In the early 1980s, its population declined by more than 80%, leading to a threatened listing under the ESA in 1993. During the 1990s, delta smelt abundance fluctuated and then increased in response to improved habitat conditions following the 1987-1992 drought. In 2002, the species' abundance again declined drastically, dropping more than 80% in just three years. In 2005, abundance of delta smelt fell to its second consecutive record low and was just 2.4% of the abundance measured when the species was listed in 1993.

Population viability and extinction risk analyses indicate that delta smelt has fallen below its "effective population size", with its present low abundance rendering the species vulnerable to inbreeding and genetic drift. The analyses, completed three years ago, predicted a 50% chance that the population would fall below its effective population size within two years, a prediction fulfilled by recent measurements of the species' record low abundance in 2004 and 2005. Other analyses using the most conservative extinction criterion predicted a 26-30% probability that the delta smelt population would fall to just 800 fish (compared to the present record low population of an estimated 25,000 fish) in the next 20 years. These high probabilities of extinction for delta smelt exceed criteria

established by the International Union for Conservation of Nature and Natural Resources for an “endangered” species.

There are multiple environmental and anthropogenic threats to the species, including: reductions in freshwater inflow to the estuary; loss of larval, juvenile and adult fish at the massive state and federal water export facilities and urban, agricultural and industrial water diversions; direct and indirect impacts of non-native species on the delta smelt’s planktonic food supply and habitat; lethal and sub-lethal effects of toxic chemicals; and, as a result of their present low population size, potential loss of the species’ genetic integrity. The recent decline of the species coincides with significant increases in Delta water exports by the state and federal water projects (seasonal water exports in the 2000s are as much as 48% higher than in the early 1990s), higher incidental take of delta smelt by the pumps, and concomitant increased alterations in internal Delta flow patterns. New analyses have demonstrated a statistically significant relationship between delta smelt population abundance and Delta export rates: high levels of water exports consistently correspond to low delta smelt abundance. Based on these analyses, current export rates are contributing to the current critically low numbers of delta smelt and are incompatible with recovery of the species. Hydrodynamic analyses of Delta flows and exports indicate that all larval and juvenile delta smelt present in the southern regions of the Delta, that part of the species’ critical habitat closest to the pumps, are likely to be entrained and lost. Ecological conditions in delta smelt habitat have also deteriorated. The invasive clam *Corbula amurensis* has reduced the abundance of the delta smelt’s zooplankton food supply. All life history stages of the species are at least periodically exposed to lethal or sub-lethal concentrations of herbicides and pesticides discharged and transported from upstream into their habitat.

State and federal management and protective regulations have proved inadequate to protect delta smelt and their estuarine ecosystem. The current export criteria in the water rights permits issued by the State Water Resources Control Board regulations allow export operations at levels that exceed those necessary to maintain healthy delta smelt populations. Dedications of water for the environment and of money for supplemental acquisitions of environmental water mandated in the 1992 Central Valley Project Improvement Act intended to reduce the negative impacts of the federal water project on fish and wildlife have not been fully or aggressively implemented. The CALFED Bay-Delta Program has been largely ineffective in addressing environmental problems in the Delta, and its future status is highly uncertain. The USFWS’s most recent Biological Opinion for protection of the species relies nearly exclusively on the CALFED Environmental Water Account, an experimental fish protection tool that, after five years of implementation, has failed to provide detectable benefits for delta smelt or any other fish species. Despite precipitously declining populations of delta smelt (and other Delta fish species), the agencies charged with protecting the species under the state and federal Endangered Species Acts are in the process of approving the South Delta Improvements Program, which would increase Delta water exports and install permanent tidal barriers that further modify Delta flow patterns and habitat.

Less than two years ago, the USFWS completed a 5-year status review for delta smelt. Based on that review, USFWS rejected a proposal to de-list the species, reaffirmed the threatened status of the species, and reported that the threats to the species had been neither eliminated nor mitigated. Since then, the condition of the species has worsened dramatically and threats to its continued existence have multiplied and escalated. The delta smelt, a potent indicator of the ecological health of the west coast's largest estuary, is at imminent risk of extinction and merits listing under the Endangered Species Act as an endangered species.

## NOTICE OF PETITION

Center for Biological Diversity  
1095 Market Street, Suite 511  
San Francisco, CA 94103  
Contact: Jeff Miller  
Phone: (415) 436-9682 x303  
Fax: (415) 436-9683  
E-mail: [jmiller@biologicaldiversity.org](mailto:jmiller@biologicaldiversity.org)

The Bay Institute  
500 Palm Drive, Suite 200  
Novato, CA 94949  
Contact: Christina Swanson, Ph.D.  
Phone: (530) 756-9021  
Fax: (530) 756-9021  
E-mail: [swanson@bay.org](mailto:swanson@bay.org)

Natural Resources Defense Council  
111 Sutter Street, 20<sup>th</sup> Floor  
San Francisco, CA 94104  
Contact: Katherine S. Poole  
Phone: (415) 875-6100  
Fax: (415) 875-6161  
E-mail: [kpoole@nrdc.org](mailto:kpoole@nrdc.org)

Petitioners Center for Biological Diversity (CBD), the Bay Institute (TBI) and Natural Resources Defense Council (NRDC) formally request that the U.S. Fish and Wildlife Service (USFWS) reclassify the listing of the delta smelt (*Hypomesus transpacificus*) from a threatened species to an endangered species under the federal Endangered Species Act (ESA), 16 U.S.C. §§1531-1544. Petitioners further request that the USFWS review whether the species warrants emergency listing, and if so, that the USFWS use its authorities to list the species as endangered on an emergency basis. This petition is filed pursuant to the authorities of 5 U.S.C. §553(e), 16 U.S.C. §1533(b)(7) and 50 C.F.R. part 424.14. The USFWS has the authority to promulgate an emergency listing rule for any species when an emergency exists that poses a significant risk to the species, under 16 U.S.C. §1533(b)(7). Such rule shall take effect immediately upon publication in the Federal Register, and shall be effective for a maximum of 240 days. This petition sets in motion a specific administrative process as defined by §1533(b)(3) and 50 C.F.R. §424.14(b), placing mandatory response requirements on the USFWS.

The delta smelt is currently designated as a threatened species under the ESA. However, the recent and continuing population decline and new information published in peer-reviewed literature, and presented at scientific meetings and CALFED Bay-Delta Program workshops described in this petition demonstrate that the species' listing should be changed to endangered – at risk of extinction. Evidence to support the listing change

presented in this petition includes: record low population abundance in the last two years; current population levels below the “effective population size” for the species cited by USFWS in their most recent 5-year review (USFWS 2004b); evidence for a significant stock recruitment relationship for the species; population viability and extinction risk analyses that indicate that the species is at high risk of extinction (Bennett 2003, 2005); and new information regarding the effects of Delta water exports on delta smelt population abundance. Due to this extreme situation, the threats documented in this petition constitute an emergency.

Because the delta smelt is primarily a freshwater fish, the USFWS has jurisdiction over this petition. A Memorandum of Understanding between the USFWS and the National Marine Fisheries Service (NMFS) signed in 1974, regarding jurisdiction over listed species, clearly places jurisdiction for the delta smelt with the USFWS (NMFS and USFWS 1974). The USFWS first proposed a threatened listing for the delta smelt on October 3, 1991 (56 FR 50075-50084) and has retained jurisdiction over the species since that time.

The Center for Biological Diversity is a nonprofit, science-based environmental advocacy organization that works to protect endangered species and wild places throughout the world through science, policy, education, citizen activism and environmental law. The Bay Institute is a non-profit organization that works to protect and restore the ecosystems of San Francisco Bay, the Sacramento-San Joaquin Delta, and the rivers, streams, and watersheds tributary to the Estuary, using a combination of scientific research, public education, and advocacy. Natural Resources Defense Council is a nonprofit, environmental organization that works to restore the integrity of the elements that sustain life – air, land and water – and to defend endangered natural places. The Center for Biological Diversity, The Bay Institute and Natural Resources Defense Council submit this petition on their own behalf and on behalf of their members and staff, with an interest in protecting the delta smelt and its habitat.

**TABLE OF CONTENTS**

Executive Summary ..... ii  
Notice of Petition ..... v  
Table of Contents .....vii

**I. NATURAL HISTORY AND STATUS OF THE DELTA SMELT ..... 1**

**A. NATURAL HISTORY ..... 1**

        1. *Description* ..... 1  
        2. *Taxonomy* ..... 1  
        3. *Distribution.* ..... 2  
        4. *Habitat Requirements.* ..... 2  
        5. *Life History* ..... 4  
        6. *Natural Mortality* ..... 7

**B. CHANGES IN DISTRIBUTION AND ABUNDANCE ..... 7**

        1. *Historic and Current Distribution.* ..... 7  
        2. *Historic and Current Abundance.* ..... 8  
        3. *Population Trend and Extinction Risk Analysis.* ..... 10

**II. CRITERIA FOR ENDANGERED SPECIES ACT LISTING ..... 12**

**A. THE DELTA SMELT IS A “SPECIES” UNDER THE ESA ..... 12**

**B. THE DELTA SMELT IS ENDANGERED UNDER THE ESA ..... 12**

        1. *Present or Threatened Destruction, Modification, or Curtailment of Its  
        Habitat or Range* ..... 13  
        2. *Overutilization for Commercial, Recreational, Scientific,  
        or Educational Purposes* ..... 19  
        3. *Disease, Competition, or Predation* ..... 20  
        4. *Inadequacy of Existing Regulatory Mechanisms* ..... 21  
        5. *Other Natural or Anthropogenic Factors* ..... 28

**III. CONCLUSION ..... 29**

**IV. SIGNATURE PAGE ..... 30**

**V. BIBLIOGRAPHY OF LITERATURE CITED ..... 31**

# I. NATURAL HISTORY AND STATUS OF THE DELTA SMELT

## A. NATURAL HISTORY

### 1. Description

The delta smelt (*Hypomesus transpacificus*) is a small (50-80 mm in length for most adults), nearly translucent, steely-blue, osmerid fish endemic to the San Francisco Bay/Sacramento-San Joaquin Delta Estuary, in California (Delta). The delta smelt was described as follows by Moyle (2002):

...slender-bodied fish that typically reach 60-70 mm SL (*standard length*; 2.36—2.76 *in*), although a few may attain 120 mm (4.73 *in*) SL. The mouth is small, with a maxilla that does not extend past the mid-point of the eye. The eyes are relatively large; the orbit width contained approximately 3.5-4 times within head length. Small pointed teeth are present on the upper and lower jaws. The first gill arch has 27-33 gill rakers and there are 7 branchiostegal rays. The gill covers lack strong concentric striations. The pectoral fins reach less than two-thirds of the way to the bases of the pelvic fins. There are 8-11 (usually 9-10) dorsal fin rays, 8 pelvic fin rays, 10-12 pectoral fin rays, and 15-19 anal fin rays. The lateral line is incomplete and has 53-60 scales along it. There are 4-5 pyloric caeca. Live fish are nearly translucent and have a steely-blue sheen to their sides. Occasionally there maybe one chromatophore (a small dark spot) between the mandibles, but usually there is none.

### 2. Taxonomy

Delta smelt is one of six species currently recognized in the *Hypomesus* genus (Saruwatari et al. 1997; as cited in Bennett 2005) and genetic analyses have confirmed that it is a well-defined species with a single intermixing population (Stanley et al. 1995; Trenham et al. 1998). Within the genus, delta smelt is most closely related to surf smelt (*H. pretiosus*), a species common along the western coast of North America. In contrast, delta smelt is a comparatively distant relation to the wakasagi (*H. nipponensis*), which was introduced into Central Valley reservoirs in 1959 and is now sympatric with delta smelt in the estuary. Delta smelt and wakasagi can also be distinguished using morphometric and pigmentation characteristics (Moyle 2002).

The U.S. Fish and Wildlife Service (USFWS) has repeatedly concluded the delta smelt is a distinct species: in an October 1991 proposed listing (56 FR 50075 50084), a March 1993 determination of threatened status (58 FR 12854 12864), a January 1994 revised critical habitat determination (59 FR 852 862), a December 1994 critical habitat determination for the species, and a 2004 5-Year Status Review for the species (USFWS 2004b).



### 3. Distribution

Delta smelt are restricted to the upper reaches of the San Francisco Bay/Sacramento-San Joaquin Delta Estuary. The species' overall distribution extends from San Pablo Bay upstream to Sacramento on the Sacramento River and Mossdale on the San Joaquin River (Radtke 1966; Moyle et al. 1992; Moyle 2002), with the actual location of the population varying seasonally and with freshwater outflow. The fish are found typically in shallow (<3 m) and/or surface waters in bays, river channels and sloughs, including in the Sacramento River, the Mokelumne River system, the Cache Slough region, the Delta, Montezuma Slough, Suisun Bay, Suisun Marsh, Carquinez Strait, the Napa River, and San Pablo Bay.

### 4. Habitat Requirements

Delta smelt are moderately euryhaline (capable of tolerating a wide range of salt water concentrations) and eurythermal (adaptable to a wide range of temperatures). The fish are found generally in brackish water (<10-12 psu [practical salinity units]), a range that corresponds well with measured salinity tolerance limits for the species (0 to 19 psu; Swanson et al. 2000). Except during the winter spawning season, most fish reside in the "low salinity zone" (0.2-2.0 psu; Bennett 2005). Delta smelt tolerate water temperatures from approximately 6°C to 25°C (depending on thermal acclimation history; Swanson et al. 2000). Temperature does not appear to control delta smelt distribution in the estuary, although temperatures >25°C may constrain delta smelt habitat during the summer and early fall (Bennett 2005).

Critical habitat for the delta smelt, designated in 1994 (59 FR 65256 65279), encompasses the areas of all water, all submerged lands below ordinary high water, and the entire water column bounded by and contained in Carquinez Strait, Suisun Bay, Goodyear, Suisun, Cutoff, First Mallard and Montezuma Sloughs, and the Sacramento-San Joaquin River Delta (as defined in Section 12220 of the California Water Code). Throughout their life cycle, the fish use different areas and habitat types within their critical habitat (and also some portions of San Pablo Bay, which is not designated as critical habitat) at different life history stages (Moyle et al. 1992; Bennett 2005; and see Life History below). For spawning, delta smelt require fresh waters as well as suitable substrate for their adhesive eggs (probably rock or submerged vegetation, although spawning microhabitat is unknown; Bennett 2005). Juveniles and sub-adults use brackish water areas in the estuary, which, depending on freshwater outflow conditions, may be located in the Delta, Suisun Bay, or San Pablo Bay.

The USFWS (1994) described the specific habitat conditions required by each life stage of delta smelt:

*Spawning Habitat* - Delta smelt adults seek shallow, fresh or slightly brackish backwater sloughs and edgewaters for spawning. To ensure egg hatching and larval viability, spawning areas also must provide suitable water quality (i.e., low concentrations of pollutants) and substrates for egg

attachment (e.g., submerged tree roots and branches and emergent vegetation). Specific areas that have been identified as important delta smelt spawning habitat include Barker, Lindsey, Cache, Prospect, Georgiana, Beaver, Hog, and Sycamore sloughs and the Sacramento River in the Delta, and tributaries of northern Suisun Bay. The spawning season varies from year to year and may start as early as December and extend until July.

*Larval and Juvenile Transport* - To ensure that delta smelt larvae are transported from the area where they are hatched to shallow, productive rearing or nursery habitat, the Sacramento and San Joaquin Rivers and their tributary channels must be protected from physical disturbance (e.g., sand and gravel mining, diking, dredging, and levee or bank protection and maintenance) and flow disruption (e.g. water diversions that result in entrainment and in-channel barriers or tidal gates). Adequate river flow is necessary to transport larvae from upstream spawning areas to rearing habitat in Suisun Bay. Additionally, river flow must be adequate to prevent interception of larval transport by the State and Federal water projects and smaller agricultural diversions in the Delta. To ensure that suitable rearing habitat is available in Suisun Bay, the 2 ppt isohaline must be located westward of the Sacramento-San Joaquin River confluence during the period when larvae or juveniles are being transported, according to the historical salinity conditions which vary according to water-year type. Reverse flows that maintain larvae upstream in deep-channel regions of low productivity and expose them to entrainment interfere with those transport requirements. Suitable water quality must be provided so that maturation is not impaired by pollutant concentrations. The specific geographic area important for larval transport is confined to waters contained within the legal boundary of the Delta, Suisun Bay, and Montezuma Slough and its tributaries. The specific season when habitat conditions identified above are important for successful larval transport varies from year to year, depending on when peak spawning occurs and on the water-year type. The Service identified situations in the biological opinion for the delta smelt (1994) where additional flows might be required in the July-August period to protect delta smelt that were present in the south and central Delta from being entrained in the State and Federal project pumps, and to avoid jeopardy to the species. The long-term biological opinion on CVP-SWP operations will identify situations where additional flows may be required after the February through June period identified by EPA for its water quality standards to protect delta smelt in the south and central Delta.

*Rearing Habitat*- Maintenance of the 2 ppt isohaline according to the historical salinity conditions described above and suitable water quality (low concentrations of pollutants) within the Estuary is necessary to provide delta smelt larvae and juveniles a shallow, protective, food-rich environment in which to mature to adulthood. This placement of the 2 ppt isohaline also serves to protect larval, juvenile, and adult delta smelt from

entrainment in the State and Federal water projects. An area extending eastward from Carquinez Strait, including Suisun Bay, Grizzly Bay, Honker Bay, Montezuma Slough and its tributary sloughs, up the Sacramento River to its confluence with Three Mile Slough, and south along the San Joaquin River including Big Break, defines the specific geographic area critical to the maintenance of suitable rearing habitat. Three-Mile Slough represents the approximate location of the most upstream extent of tidal excursion when the historical salinity conditions described above are implemented. Protection of rearing habitat conditions may be required from the beginning of February through the summer.

*Adult Migration* - Adult delta smelt must be provided unrestricted access to suitable spawning habitat in a period that may extend from December to July. Adequate flow and suitable water quality may need to be maintained to attract migrating adults in the Sacramento and San Joaquin River channels and their associated tributaries, including Cache and Montezuma sloughs and their tributaries. These areas also should be protected from physical disturbance and flow disruption during migratory periods.


## **5. Life History**

The life history of delta smelt has been extensively described by Moyle et al. (1992), Moyle (2002), Bennett (2005), and the most recent Biological Opinion for the species (USFWS 2004a). Briefly, delta smelt is an annual species although there is evidence that, at least in some years, a small percentage of the population (<10%) survives to two years and may spawn in one or both years (Bennett 2005). Compared to other annual species, delta smelt exhibit relatively low fecundity with most females producing fewer than 3,000 eggs (Bennett 2005; Mager 1996). Spawning occurs in fresh water during the winter and spring (February-June; peak spawning April-May) and at water temperatures between 12°C and 18°C. Hatching occurs approximately 8-14 days later depending on water temperature. Newly hatched larvae are small (4.5-6 mm) and, after yolk-sac absorption and development of the jaw and mouth parts, they begin feeding on unicellular algae, rotifers, and/or sub-adult copepods. As the fish grow, their diet shifts nearly exclusively to copepods. Larval and juvenile fish move and/or are transported by flow and tides downstream from freshwater areas in the upper Delta (and, in some years, the Napa River and upper Suisun Marsh) to the brackish low-salinity zone where they rear for the summer and fall. In late fall and early winter, maturing adults begin their diffuse upstream migration to freshwater spawning areas.

Delta smelt exhibit a weak but significant stock recruitment relationship. Using data from California Department of Fish and Game (CDFG) surveys that measure the abundance of juvenile and sub-adult/adult delta smelt (see Historic and Current Abundance, below), correlation and regression analyses show that the abundance of juvenile fish is significantly and directly related to the abundance of the adults that produced them (Figure 1 and Equation 1).<sup>1</sup>

---

<sup>1</sup> Population abundance of delta smelt is monitored by two California Department of Fish and Game surveys. The Summer Tow Net Survey (TNS), conducted during June and July, measures the abundance of



**Figure 1.** The relationship between the abundance of juvenile delta smelt (log TN Index) and the abundance of the adult fish that produced them (log FMWT Index for the previous year). Data are for 1969 to 2005. Regression equation and associated statistics are shown with the graph and are provided in the text. Data source: California Department of Fish and Game.

$$\text{Log TNS Index} = -1.41 + 0.83(\text{log FMWT Index previous year}) \quad \text{(Equation 1)}$$

n=35; p<0.001; r<sup>2</sup>=0.341; standard error of the estimate (SEE)=0.437  
 where: log TNS Index is abundance of juvenile delta smelt; and  
 log FMWT Index is the abundance of sub-adult/adult delta smelt.


Similarly, the abundance of adult fish measured the following fall is significantly and directly related to the abundance of the population as juveniles measured earlier in the year (Figure 2 and Equation 2).

$$\text{Log FMWT Index} = 2.21 + 0.46(\text{log TNS Index}) \quad \text{(Equation 2)}$$

n=35; p<0.001; r<sup>2</sup>=0.337; SEE=0.347

---

juvenile delta smelt. The Fall Mid Water Trawl (FMWT), conducted from September to December, surveys sub-adult and adult delta smelt. From each of these surveys, delta smelt population abundance is calculated as an annual abundance “index”. Because the FMWT and TNS abundance index data are not normally distributed, for these and other statistical analyses, the indices were log<sub>(10)</sub> transformed (similar to the approach used by Bennett 2005).



**Figure 2.** The relationship between the abundance of adult delta smelt (log FMWT Index) and the abundance of juvenile measured earlier in the same year (log TNS Index). Data are for 1969 to 2005. Regression equation and associated statistics are shown with the graph and are provided in the text. Data source: California Department of Fish and Game.

There is considerable disagreement among experts regarding the possible influence of density dependence (i.e., regulation of the size of the population by mechanisms that are themselves controlled by the size of that population, such as availability of food resources) on delta smelt population dynamics and abundance. Bennett (2003, 2005) suggested that the abundance of adult fish might be limited by density dependent effects on juveniles. The CDFG (2003) disagreed in their comment letter to the USFWS for the recent 5-year status review for delta smelt (USFWS 2004b), citing lack of credible mechanism(s), the recent low abundance of delta smelt, and the weakness of the statistical evidence for density dependence. The fact that delta smelt are not currently abundant relative to their recent past abundance or relative to the other species living in the Delta suggests that the species has not exceeded the carrying capacity of its habitat and that its abundance is not limited by density dependent effects. Regardless, Bennett's (2005) analysis indicated that, since the early 1980s, density dependent effects did not occur at population abundance levels lower than 400-450 (as measured by the Fall Mid Water Trawl Abundance Index; and see Historic and Current Abundance below), levels that are roughly comparable to or somewhat lower than the recovery criteria identified for the species (Moyle et al.1996).

## 6. Natural Mortality

The most recent review of delta smelt status reports that there is insufficient evidence for effects of disease, competition or predation on delta smelt population abundance (USFWS 2004b). However, diseases and parasites of delta smelt have not been well studied and numerous introduced species are abundant in the estuary. Competition and predation by introduced species has been shown to affect delta smelt size and condition (Souza et al. 2005; Teh 2005) but its role in affecting delta smelt population abundance has not been demonstrated (see discussion of Disease, Competition, or Predation below).

### B. CHANGES IN DISTRIBUTION AND ABUNDANCE

#### 1. Historic and Current Distribution

**Historic distribution:** The delta smelt's historic distribution extended from San Pablo Bay upstream to Sacramento on the Sacramento River and Mossdale on the San Joaquin River (Radtke 1966; Moyle et al. 1992; Moyle 2002), with the actual location of the population varying seasonally and with freshwater outflow. Based on results of CDFG sampling surveys for juveniles during the Summer Townet Survey (TNS) and adults during the Fall Mid Water Trawl (FMWT) survey conducted since the late 1950s as well as earlier published reports (cited in the references above), delta smelt were found in shallow (<3 m) and/or surface waters in bays, river channels and sloughs, including in San Pablo Bay, Carquinez Strait, the Napa River, Suisun Bay, Suisun Marsh, Montezuma Slough, the Sacramento River, the Cache Slough region, the Mokelumne River system, the San Joaquin River, Old and Middle Rivers, and all other channels and sloughs in the Delta with accessible subtidal, shallow-water habitat. In general, the rivers, channels, and sloughs in the Delta function as both rearing habitat for larval and juvenile delta smelt and a migration corridor for both the adult and juvenile life stages while the brackish, open bay areas serve as rearing habitat for older juveniles and sub-adult delta smelt (see Habitat Requirements and Life History).

**Current distribution:** Since the late 1970s, large areas of historic delta smelt habitat (and designated critical habitat) within the Delta have been rendered unsuitable for a key life history stage of the species: the distribution of the species has been substantially reduced. Based on results from the summer TNS, which has been conducted annually since 1959, juvenile delta smelt have virtually disappeared from the Delta in areas south of the San Joaquin River (Miller 2000; CDFG 2003; Fleming and Nobriga 2004). Most of the environmental characteristics of these south Delta channels and sloughs known to be necessary for the species (e.g., temperature, salinity, water depth) have not changed. Recent surveys for adult delta smelt (CDFG Spring Kodiak Trawl) and larvae (CDFG 20 mm Survey) indicate that this region of the Delta is still used for spawning in the spring. But, by two to three months later, few or no juvenile delta smelt are collected during the summer TNS, a marked change compared to results of the first 20 years of the survey. This is the region of the Delta most directly affected by State Water Project (SWP) and Central Valley Project (CVP) water export operations and by the installation and spring-summer operations of three south Delta tidal gate barriers designed to maintain water

levels and water quality for local agricultural water diverters. The period during which juvenile delta smelt disappeared from this portion of their historic habitat (i.e., the late 1970s) coincides with substantial increases in the amounts of water exported, increases in frequency of high volume export operations, and the initiation and increasing duration of south Delta barrier operations (Simi and Ruhl 2005; Fleming and Nobriga 2004; and see Figure 4).

## 2. Historic and Current Abundance

More than a dozen regular surveys collect data on delta smelt within the estuary (USFWS 2004b) but the population abundance estimates used for most analyses and for tracking the species' population status are calculated from results of two surveys conducted by the CDFG: the TNS which samples juvenile delta smelt and the FMWT which samples sub-adult and adult delta smelt. From each survey, an abundance index is calculated by extrapolating the numbers of fish caught at 30 to 80 fixed stations, using a weighting factor that accounts for differences in water volume at various sub-regions from San Pablo Bay through the Delta (Bennett 2005). A subset of stations and months from the FMWT is also used to calculate the Recovery Index (Moyle et al. 1996). None of these indexes provide direct measurements of actual population abundance. However, total population size calculated from the raw survey data and habitat volume estimates by Bennett (2005) show a good linear correspondence with the abundance indices, suggesting that the indices represent reasonable estimates of relative population abundance.


**Historic abundance:** As recently as 30 years ago, delta smelt was one of the most common and abundant of the pelagic fishes caught in the estuary, as indicated by its abundance in CDFG trawl catches (Erkkila et al. 1950; Radtke 1966; Stevens and Miller 1983). For the first 16 years of the FMWT survey (1967-1982), the most comprehensive survey for the species, the abundance index averaged 806, a value that roughly corresponds to an estimated 800,000 adult fish (Bennett 2005). But, in the early 1980s, the population declined by more than 80% (based on the FMWT Index; the decline was 95% based on the TNS Index; Table 1 and Figure 3). According to Bennett (2005) this precipitous decline was the result of the combination of extremely high winter and spring outflows, elevated springtime water temperatures and resultant short duration spawning periods, and two consecutive years of extremely high levels of incidental take at the SWP and CVP water export facilities. From 1983 through 1992, delta smelt abundance (measured as the FMWT Index) was consistently low, averaging just 33% of the 1967-1982 average abundance levels. In 1993, abundance increased considerably and, throughout the rest of the decade, population levels fluctuated but were, on average, roughly double those measured during the 1983-1992 period (Table 1). This increase prompted several stakeholder groups to challenge the listing, an action that ultimately led to the recent 5-year status review for the species (USFWS 2004b).

**Table 1.** Abundance of delta smelt as measured by the CDFG Summer Tow Net Survey (TNS, juvenile delta smelt), Fall Mid Water Trawl Survey (FMWT, adult delta smelt), and the Recovery Index (based on FMWT data, adult delta smelt).

Year	TNS Index	FMWT Index	FMWT averages	Recovery Index	Comments
1959	12.1				Comparisons are based on FMWT Indices
1960	25.4				
1961	21.3				
1962	24.9				
1963	1.8				
1964	24.6				
1965	6.0				
1966	no data				
1967	no data	414	1967-1982: 806	139	
1968	no data	696		251	
1969	2.5	315		128	
1970	32.5	1673		598	
1971	12.5	1303		352	
1972	11.1	1265		551	
1973	21.3	1145		305	
1974	13.0	no data		no data	
1975	12.2	697		239	
1976	50.6	360		22	
1977	25.8	481		146	
1978	62.5	572		108	
1979	13.3	no data		no data	
1980	15.8	1653		312	
1981	19.8	374		78	
1982	10.7	330		37	
1983	2.9	132	1983-1992: 267	17	Population declines 84% (from 1967-1982 FMWT).
1984	1.2	182		51	
1985	0.9	110		29	
1986	7.9	212		70	
1987	1.4	280		72	
1988	1.2	174		67	
1989	2.2	366		76	
1990	2.2	364		81	
1991	2.0	689		171	
1992	2.6	156		26	
1993	8.2	1078	1993-2001: 572	400	Delta smelt listed as threatened
1994	13	102		19	Abundance is 9% of 1993 level.
1995	3.2	899		252	Abundance is 83% of 1993 level.
1996	11.1	127		28	Abundance is 12% of 1993 level.
1997	4.0	303		62	Abundance is 28% of 1993 level.
1998	3.3	420		169	Abundance is 39% of 1993 level.
1999	11.9	864		322	Abundance is 80% of 1993 level.
2000	8	756		265	Abundance is 70% of 1993 level.
2001	3.5	603	1999-2001: 741	314	Abundance is 56% of 1993 level.
2002	4.7	139		33	Population declines 81% (from 1999-2001 FMWT). Abundance is 13% of 1993 level.
2003	1.6	210		101	Recovery criteria met (this year only). Abundance is 19% of 1993 level.
2004	2.9	74		25	Abundance is 7% of 1993 level.
2005	0.3	26		4	Abundance is 3% of 1967-1982 average, 4% of 1999-2001 average, and 2% of 1993 level.



**Current abundance:** In 2002, the delta smelt population again experienced a drastic population decline, dropping more than 80% from levels measured during the previous three years (1999-2001 average) (based on the FMWT Index; the decline was 77% based on the 2003 TNS Index compared to the 1999-2002 average; Table 1 and Figure 3). Since then, the population has continued to decline, falling to what was then a record low in 2004 (FMWT=74, or an estimated 60,000-70,000 adult fish) and then plummeting again by 65% to a FMWT Index of just 26 in 2005, the lowest abundance level ever recorded for sub-adult and adult delta smelt. Based on Bennett's (2005) analyses, the 2005 FMWT Index roughly corresponds to fewer than 25,000 adult fish.



**Figure 3.** Trends in abundance of delta smelt as measured by two different surveys. The upper panel shows the abundance index calculated from the Summer Townet Survey (TNS, 1959-2005), which samples juvenile delta smelt. The lower panel shows the abundance Index calculated from the Fall Midwater Trawl Survey (FMWT, 1967-2005), which samples sub-adult and adult delta smelt. Data source: California Department of Fish and Game.

### 3. Population Trend and Extinction Risk Analysis

By the time the delta smelt was listed as threatened in 1993, the species had experienced a decade of chronically low population abundance followed by the beginnings of an apparent recovery, with both juvenile and adult abundance indices increasing (Table 1 and Figure 3). In 1993, delta smelt abundance was the highest measured since the major population decline in the early 1980s. During the rest of the 1990s, abundance fluctuated but generally continued to improve. This period corresponded with favorable water

temperature and flow conditions. The most recent population decline began in 2000 and, in 2002, abundance of adult delta smelt dropped by more than 75% compared to the previous year. The following year, abundance of the juvenile fish was 66% lower than that measured in the previous year. Since then, population abundance as measured by both life-stage specific surveys has continued to decline. By 2005, abundance of juvenile delta smelt was just 3.7% of that measured in 1993 when the species was listed under the ESA and the abundance of adult delta smelt was just 2.4% of the 1993 level.

Using data on past measured population abundance and variability, Bennett (2003, 2005) conducted a population viability analysis for delta smelt, calculating probabilities and predicted time frames for extinction of the species.<sup>2</sup> In the first step of his analysis, Bennett (2003) estimated the delta smelt population would be below its “effective population size”, the population level below which a species is subject to inbreeding and genetic drift, if the FMWT Index fell below 100 for two consecutive years. At the time of this analysis, this abundance level was similar to the lowest FMWT Index ever measured for the species (1994 FMWT Index of 102, or an estimated 86,203 adult fish). The USFWS 5-year status review (2004b) recognized this definition of an effective population size and noted that this population size reflected an “unprecedented low number of delta smelt.” Based on this analysis, the current delta smelt population is well below its effective population size, with the 2004 and 2005 FMWT Index values of 74 and 26 respectively (Table 1).

Using the now defined effective population size as a benchmark, Bennett (2003, 2005) then conducted an extinction risk analysis using three progressively more extreme population size criteria to simulate extinction: 80,000, 8,000, and 800 fish. For each population level, the analysis used FMWT data for the 1982-2003 period to calculate the median time to a 50% probability of extinction and the probability that extinction would occur within 20 years. For the highest population level, the analysis predicted that the median time to when there was a 50% chance that the population would fall below 80,000 fish was only 1.2 to 1.5 years, a prediction confirmed one year later when the FMWT Index fell to 74. Using the lower extinction levels, Bennett (2005) predicted that the median time to 50% probability of extinction defined as a population size of just 8,000 fish was only 20 years and, for a population size of 800 fish, 42 to 55 years. For this lowest extinction level (800 fish), there was a 26-30% probability that extinction would occur within 20 years. Bennett (2005) compared his results to criteria developed by the International Union for Conservation of Nature and Natural Resources, which defines an endangered species as one with a 20% probability of extinction in 20 years, and stated that “delta smelt qualifies for Endangered ... status” (page 55).

---

<sup>2</sup> Another extinction probability analysis conducted by the San Luis & Delta-Mendota Water Authority (2002) was considered by the USFWS in their 5-year review (USFWS 2004b). However, independent peer-review of this analysis by USGS (2003) concluded that this extinction probability analysis and estimate was seriously flawed.

## **II. CRITERIA FOR ENDANGERED SPECIES ACT LISTING**

### **A. THE DELTA SMELT IS A “SPECIES” UNDER THE ESA**

The Delta smelt is indisputably a species, as discussed in the section on taxonomy above. The USFWS has acknowledged the delta smelt is a distinct species in the October 1991 proposed listing (56 FR 50075 50084), a March 1993 determination of threatened status (58 FR 12854 12864), a January 1994 revised critical habitat determination (59 FR 852 862), a December 1994 critical habitat determination for the species, and a 2004 5-year status review for the species (USFWS 2004b).

### **B. THE DELTA SMELT IS ENDANGERED UNDER THE ESA**

The ESA defines an “endangered” species as “any species which is in danger of extinction throughout all or a significant portion of its range...” and a “threatened” species as “any species which is likely to become an endangered species within the foreseeable future throughout all or a significant portion of its range.” The threats to a species’ survival may be categorized according to the ESA as:

- (A) the present, or threatened destruction, modification, or curtailment of its habitat or range;
- (B) over-utilization for commercial, recreational, or educational purposes;
- (C) disease or predation;
- (D) inadequacy of current regulatory mechanisms; or
- (E) other natural or manmade factors affecting its continued existence.

According to the USFWS 5-year status review (USFWS 2004b), there are sufficient data and analyses available to identify (A) habitat destruction, modification and curtailment; (D) inadequacy of current regulatory mechanisms; and (E) other natural and manmade factors as present and ongoing threats to the species. The USFWS (2004b) also concluded that for delta smelt, “most threats which were discussed in the original listing remain.” In this petition to list delta smelt as endangered we concur with the threats analysis published in the 5-year status review (USFWS 2004b) and add new analyses and information to further describe the nature, degree, and increasing severity of the dual threats of habitat modification and the inadequacy of existing regulatory mechanisms to protect the species.

Delta smelt are environmentally sensitive because they have a short life span, have a limited diet, have a low fecundity for a fish producing planktonic larvae, are poor swimmers, are easily stressed, and reside primarily in the interface between salt and freshwater within a limited geographic extent. The species is highly vulnerable to extinction because of its short life span, present small population size, and restricted distribution. As noted by Moyle (2002), a substantial population is necessary to keep delta smelt from becoming extinct, and the species has recently fallen below the effective population size for viability (as defined by Bennett 2003) to an “unprecedented low number of delta smelt” (USFWS 2004b). Multiple factors are thought to be contributing to the continuing population decline of the species, including: reductions in freshwater

outflows to the estuary; direct and indirect adverse impacts of Delta water diversions and exports; reductions in abundance of prey food organisms; lethal, sub-lethal and indirect effects of toxic substances; disease, competition, and predation; and loss of genetic integrity. The magnitude and frequency of occurrence of many of these factors are increasing.

The delta smelt indisputably merits immediate emergency listing under the ESA as an endangered species. Furthermore, the listing decision must be made solely on the basis of the best scientific and commercial data available, under 16 U.S.C. §1533(b)(1)(A). The legislative history of this provision clearly states the intent of Congress to “ensure” that listing decisions are “based solely on biological criteria and to prevent nonbiological criteria from affecting such decisions” (H.R. Rep. No. 97-835, 97th Cong. 2d Sess. 19 (1982)). As further stated in the legislative history, “economic considerations have no relevance to determinations regarding the status of species.” Therefore, political and economic arguments may not be considered by the USFWS in its determination of whether to list this species. The factors threatening the continued survival of the delta smelt are detailed below.

### **1. Present or Threatened Destruction, Modification, or Curtailment of Its Habitat or Range**

Since delta smelt live entirely within the estuary, confined to a relatively small geographic range of low salinity habitat, changes in Delta water quality, water inflow and export regimes, and the estuarine food web have direct and significant impacts on the species.

#### Water Exports and Diversions

Several recent analyses have demonstrated that Delta water exports are having significant negative effects on delta smelt population abundance and suggest that the adverse effects of water exports on delta smelt may have increased since the listing of the species as threatened.

The Delta, the heart of delta smelt spawning and early rearing habitat, is a major hub for California’s water management system and a region of intensive irrigated agriculture. Water management operations of the massive SWP and the federal CVP in the Delta and its Sacramento and San Joaquin River watersheds have had major and often detrimental effects on freshwater inflow to the Delta, in-Delta hydrodynamics, and water quality. For example, during periods of low freshwater inflow and high volume water exports, conditions that are common during periods when the delta smelt population is distributed in the Delta, as much as 65% of the total freshwater inflow may be diverted, and net movement of water and entrained plankton, larvae and small fish in the central and southern Delta is towards the export facilities. At times, the net flow of the San Joaquin River is reversed, confusing and delaying up-migrating adult fish and impairing downstream transport of larval and juvenile delta smelt from the upper estuary where they were spawned to their brackish water rearing habitat. The fish screens at the SWP

and CVP fish facilities are known to be inadequate for protection of delta smelt (as well as for most other species; e.g., Bowen et al. 2004). Despite evidence of the limited effectiveness of these facilities for fish protection, and evidence that this effectiveness is further deteriorating (Bowen et al. 2004), plans to upgrade or replace the facilities have been delayed indefinitely (SDFFF 2005).

In addition to the SWP and CVP water export facilities, more than 1,800 smaller diversions extract water for local consumptive use, and two power plants use Delta water for cooling (Herren and Kawasaki 2001). The USFWS (2004b) concluded that the power plants at Pittsburg and Antioch with flow-through cooling, within the region of seasonal maximum abundance of delta smelt, can impose significant mortality on delta smelt. The vast majority of the other Delta diversions are unscreened (Herren and Kawasaki 2001).


Water exports and diversions adversely affect delta smelt directly by entrainment (i.e., movement of fish due to the hydraulic effects of pumping and lethal removal of the fish from its habitat through unscreened or inadequately screened water diversions) or impingement (getting stuck on fish or debris screens at pumps); and indirectly by adversely modifying their critical habitat (e.g., by altering in-Delta hydrodynamics, reducing freshwater outflow and shifting the location of suitable brackish water rearing habitat, and/or removing planktonic food organisms). The species is at greatest risk from water diversions at two critical times in its life cycle, first as larvae and juveniles when the young fish move downstream to brackish rearing habitat (spring and early summer) and then again as pre-spawning and spawning adults when the fish move up into the Delta for reproduction (winter and early spring). All of these life stages are known to be lethally entrained at government, urban, industrial, and agricultural diversions in the Delta (e.g., Matica and Sommer 2005; Nobriga et al. 2002; Moyle 2002). Entrainment loss of delta smelt >20 mm in length is monitored and reported as incidental take at only the SWP and CVP water export facilities.<sup>3</sup> Incidental take of delta smelt at the other diversions and of larval and juvenile fish <20 mm in length at the government water project facilities is neither monitored nor reported and no effort is made to rescue these fish. Therefore, the incomplete data on incidental take are of limited value for evaluating the effects of water export activities on delta smelt population levels, although they do provide useful information on the timing and presence of the fish in the south and central Delta.

Since the delta smelt was first listed as threatened under the ESA in 1993, annual water exports from the Delta have increased by 20% (1994-1998 average: 4.773 million acre-

---

<sup>3</sup> Incidental take reported for delta smelt at the state and federal water export facilities is known to be a gross underestimate of the actual numbers of fish lethally entrained. The daily take number is calculated from counts of fish >20 mm in length collected in regular sub-samples of water bypassed through two sequential sets of louvers and/or fish screens. However, efficiency of the louvers to remove delta smelt from the diverted water is known to be low for delta smelt (i.e., under many conditions, more than half of the diverted fish pass through the louvers and are transported uncounted to the pumps; Bowen et al. 2004). Delta smelt larvae and juveniles <20 mm in length are not counted, but the loss of larval and juvenile delta smelt to entrainment is estimated at several million each year (USFWS 2004b). In addition, unknown proportions of the fish entrained into the facilities are lost to predation and/or other mortality factors and never reach the fish salvage facilities to be counted.

feet [MAF] per year; 2000-2004 average: 5.719 MAF per year).<sup>4</sup> This increase continues a long-term trend of increases in Delta water exports (Figure 4, top panel). During the periods when delta smelt are vulnerable to entrainment, the relative increases in exports have been even greater, with a 48% increase in winter exports (December-March, when pre-spawning and spawning adults are in the Delta, Figure 4, middle panel) and a 28% increase in spring exports (March-July, when larvae and small juveniles are in the Delta, Figure 4, bottom panel).




**Figure 4.** Combined water exports (million acre-feet, MAF) of the Central Valley Project and the State Water Project from 1967-2004. The upper panel shows total export volumes for each water year. The middle panel shows exports for the winter period (December-March). The lower panel shows springtime exports (March-July). Data source: California Department of Water Resources, Dayflow.

#### Direct Impacts:

The CDFG (2003) expressed concern that entrainment of delta smelt at the CVP and SWP could be a major source of population impacts under certain conditions. Data from the summer TNS show an almost complete disappearance of juvenile delta smelt in south

<sup>4</sup> Exports are calculated as the combined annual total (using the water year calendar) for Central Valley Project and State Water Project exports using data from Dayflow (California Department of Water Resources).

Delta sampling stations by the mid-1970s, coincident with a trend of increasing combined water exports from the south Delta (CDFG 2003). CDFG (2003) estimated losses of delta smelt juveniles to SWP and CVP operations to range from 11 to 46% of the population annually.



**Figure 5.** Recent patterns in incidental take (or salvage) of delta smelt during the November-March period. Top panel: total salvage (# fish); Middle panel: salvage density (# of fish/thousand acre-feet); and Bottom panel: salvage density in relation to preceding FMWT abundance Index (salvage density/FMWT Index previous year). Source: Herbold et al. 2005, Figure 3.

Analyses conducted by Herbold et al. (2005) as part of the ongoing multi-agency research program to investigate the recent pelagic fish declines in the Delta indicate that the direct impacts of water exports on delta smelt (and other Delta pelagic fish species) during the winter have increased in recent years, coincident with substantial population declines measured for all the affected species.<sup>5</sup> Beginning in 2000, incidental take (also referred to as “salvage”) of adult delta smelt increased markedly (Figure 5, top panel), concurrent with substantial increases in exports during the same period (see Figure 4, middle panel). In 2003, one year after the most recent delta smelt population decline, direct loss of adult

<sup>5</sup> Herbold et al. (2005) used the November-March period for their analyses.

delta smelt at the pumps in relation to the species' population abundance reached its highest level in more than ten years (Figure 5, bottom panel). On the basis of these and other analyses conducted by researchers investigating the pelagic organism decline, the "Winter Adult Entrainment Hypothesis" is one of two leading hypotheses to explain the pelagic fish decline that will be the focus of continuing research in 2006 (Armor et al. 2005).<sup>6</sup>

The USFWS (2004b) noted that the California Department of Water Resources (CDWR) and the U.S. Bureau of Reclamation (BOR) proposal to replace temporary barriers at the SWP and CVP with permanent barriers could result in additional effects to delta smelt. The current temporary barriers physically prevent delta smelt movement and change south Delta hydrodynamics, increasing central Delta flows toward the state and federal water export facilities (USFWS 2004b). The proposed permanent barriers operations may include barrier closures during additional periods and may include different operations that could affect delta smelt (USFWS 2004b). When delta smelt occur in areas influenced by the barriers, entrainment losses at the state and federal export facilities could increase (USFWS 2004b).

#### Population Level Impacts:

Recent analyses have shown that delta smelt population abundance is significantly and inversely related to seasonal export rates (Swanson 2005).<sup>7</sup> Figure 6 and Equations 3 and 4 show that export rates during the winter (December-March) significantly affect the abundances of adult and juvenile delta smelt measured later that year; for both life stages, abundance decreased with increases in exports.<sup>8</sup>

Adult delta smelt (1967-2004):

$$\text{Log FMWT} = 3.042 - 0.297(\text{Dec-Mar exports, MAF}) \quad \text{(Equation 3)}$$

n=36; p=0.003; r<sup>2</sup>=0.0239, SEE=0.326

Juvenile delta smelt (1967-2004):

$$\text{Log TNS} = 1.336 - 0.347 (\text{Dec-Mar exports, MAF}) \quad \text{(Equation 4)}$$

n=36 p=0.017, r<sup>2</sup>=0.155, SEE=0.450

Analyses for the alternative winter periods (November-February, November-March, and December-February) yielded similar and statistically significant results (although p values were somewhat higher and r<sup>2</sup> values were somewhat lower).


---

<sup>6</sup> In 2005 the Interagency Ecological Program began an intensive research program to investigate the cause(s) of the Pelagic Organism Decline (POD) in the Delta. Preliminary results of this research are available at: [http://science.calwater.ca.gov/workshop/workshop\\_pod.shtml](http://science.calwater.ca.gov/workshop/workshop_pod.shtml).

<sup>7</sup> These analyses were conducted by C. Swanson and earlier versions of these results were presented at the CALFED Bay-Delta Program-sponsored Environmental Water Account Review Workshop, Dec. 7-8, 2005, Sacramento, CA. Presentation and supplemental report are available at: <http://science.calwater.ca.gov/workshop/ewa.shtml>.

<sup>8</sup> For reference, export of 2.0 MAF during the December-March period corresponds to an average daily combined export rate of the SWP and the CVP of approximately 8,300 cubic feet per second (cfs).





**Figure 6.** The relationship between winter (December-March) export amounts and subsequent abundance of delta smelt. a) sub-adult and adult delta smelt as measured by the FMWT Index (using data from 1967-2004); and b) juvenile delta smelt as measured by the TNS Index (using data from 1969-2004). For each graph, the regression, 95% confidence limits and the prediction limits are shown calculated for the entire datasets. The open symbols and the dark gray regression line highlight the years since the delta smelt was listed under the ESA (1994-2004). Data Sources: California Department of Fish and Game, California Department of Water Resources, Dayflow.

Large scale ecological changes have occurred in the Delta during the past 30 years, such as the establishment of the invasive clam *Corbula amurensis* and its impacts on the planktonic food web, but they do not strongly affect the results of these types of correlation and regression analyses. For example, the significant relationship between winter exports and the subsequent population abundance of adult delta smelt was apparent in the 20 years prior to the clam's invasion (1967-1985, Equation 5).

Adult delta smelt (1967-1986):

$$\text{Log FMWT} = 3.109 - 0.353(\text{Dec-Mar exports, MAF}) \quad \text{(Equation 5)}$$

n=18; p=0.013; r<sup>2</sup>=0.0329, SEE=0.308

Linear regression using smaller subsets of more recent years (e.g., post-*Corbula* invasion, 1987-2004 or 2005; post-ESA listing, 1994-2004 or 2005) were not statistically significant but both the slopes and intercepts of the relationships were very similar to those generated using the entire dataset (e.g., 1994-2004(5): open symbols and grey

regression line in Figure 6). The significant relationship between winter exports and abundance was not “driven” by the low abundances measured during the past three or four years. For example, after excluding the three most recent years for the FMWT abundance indices (2002-2004) from the dataset, the regression was still significant ( $p=0.02$ ) and the slope and intercept were similar to those generated with the entire dataset. Given that the significant relationship between winter exports and adult abundance was detectable by 2002 (and before), this indicates that the low abundances measured during the past three years, a period during which winter exports were at near record high levels, were predictable as early as three years ago.

The abundance of juvenile delta smelt was also significantly affected by spring-summer exports (March-July). The linear regression for this relationship is:

$$\text{Log TNS} = 1.429 - 0.369(\text{Mar-July exports, MAF}) \quad \text{(Equation 6)}$$

$N=36; p=0.047; r^2=0.111; \text{SEE}=0.462$

In 1993, the USFWS (1993) identified 21 major federal, state, local or private organization proposals for increased exports. Since that time, Delta water exports and corresponding impacts on delta smelt have increased and they are projected to continue to increase in the future. The recent 5-year review (USFWS 2004b) noted that the potential threat of increased demands on surface water resources in the Central Valley and Delta was growing, citing planned or proposed new water diversion projects such as the Freeport Regional Water Project, increases in pumping capacity at the SWP pumping plant as part of the South Delta Improvement Project, the California Aqueduct/Delta-Mendota Canal inter-tie to allow increased pumping at the CVP pumping plant, Empire Tract on the San Joaquin River; and potential expanded water storage capacity projects at Los Vaqueros, north of the Delta off-stream storage, Shasta Reservoir, in-Delta storage, and south of the Delta surface and groundwater storage projects. The USFWS (2004b) concluded that the increased storage and diversion capacity would likely result in lower freshwater outflows to the estuary, higher water exports from the Delta, and greater entrainment of delta smelt.

## **2. Overutilization for Commercial, Recreational, Scientific, or Educational Purposes**

Over-utilization for commercial, recreational, scientific, or educational purposes is not known or thought to be a factor in the decline of the delta smelt population (USFWS 2004b). Delta smelt may be harvested as a non-target by-catch in commercial bait fisheries for other baitfish species and some scientific collecting is conducted for the delta smelt, but the USFWS (1993) did not believe these activities were adversely affecting the species.

### 3. Disease, Competition, or Predation

There is no evidence that disease, competition, or predation has caused the delta smelt population to decline (Moyle et al. 1996, 2004b), although diseases and parasites of delta smelt have been little studied. Introduced species are abundant in the estuary, including non-native invertebrates and fishes that feed on phytoplankton, zooplankton and/or small fish, and may adversely affect the delta smelt (USFWS 2004b). There is some evidence that increased predation pressure on the planktonic food web by introduced species such as the invasive clam *Corbula amurensis* have reduced food availability for delta smelt (and other planktivorous fishes in the estuary; Kimmerer and Orsi 1996; Armor et al. 2005) and recent studies have related reduced plankton biomass to reduced delta smelt size (Souza et al. 2005) and liver glycogen levels (Teh 2005), although growth rates are unchanged (Souza et al 2005). However, to date, there is no evidence that competition for food or space with other aquatic organisms has affected delta smelt population abundance.

Introduced striped bass may have caused an increase in predation on all size classes of the delta smelt (USFWS 2004b). Efforts to enhance striped bass populations by planting large numbers of juveniles from hatcheries could have had a negative effect on other pelagic fishes in the estuary once the striped bass reached a size where they begin preying upon fish. The enhanced predator populations, without a concomitant enhancement of prey populations such as delta smelt, may have resulted in excessive predation pressure on prey species. Striped bass appear to have switched to piscivorous feeding habits at smaller sizes than they did historically following severe declines in abundance of mysid shrimp (CDWR 2003). The CDFG has completed a Habitat Conservation Plan for their striped bass management program which includes measures designed to help conserve delta smelt.

Introduced planktivores (fish that eat plankton) such as threadfin shad (*Dorosoma pretense*) and inland silverside (*Menidia beryllina*) compete for the zooplankton food of delta smelt, or alter the species composition of the zooplankton community, thereby further decreasing the ability of the delta smelt population to recover. Although the delta smelt has managed to coexist with these species in the past, it is possible that, at low population levels, interactions with them could prevent recovery. In particular, inland silversides are frequently collected in areas where delta smelt may spawn and they could be major predators on eggs and larvae (Bennett 1995, 2005; CALFED 2001). Estimates of abundance of delta smelt and silversides are negatively correlated, suggesting that inland silversides may be an important predator on larval delta smelt and a competitor for copepod prey. Silversides often occur in dense schools near shorelines and their occurrence may detract from the value of shallow water habitat created to aid delta smelt restoration. Since the early 1980s, there have been increases in other potential larval fish predators such as tagged Chinook salmon smolts released in the delta for survival experiments and non-native centrarchids (CDWR 2003). Introduced species such as the chameleon goby (*Tridentiger trigonocephalus*) and yellowfin goby (*Acanthogobius flavimanus*) may prey on delta smelt eggs and larvae and interfere with recovery of the species.

#### **4. Inadequacy of Existing Regulatory Mechanisms**

The recent delta smelt population trends, substantial increases in environmental and anthropogenic stressors already identified as threats to the species (i.e., Delta water exports), concurrent increases in incidental take, and statistical analyses that reveal significant relationships between those stressors and the recent and long-term declines in the delta smelt population clearly show that current regulatory mechanisms do not provide adequate protection for the species. In addition, specific protections and management tools identified in the 2004 Biological Opinion (USFWS 2004a) have proved inadequate and, apparently, unavailable to state and federal fish agency managers attempting to use them to protect the species. Current regulatory mechanisms and their inadequacy to prevent the imminent extinction of the delta smelt are discussed below.

##### State Threatened Listing:

The delta smelt was listed by the State of California as a state threatened species under the California Endangered Species Act (CESA) in December 1993. While state listing theoretically provides some minimal measure of protection to the species because state agencies are required to consult with the CDFG if any project they fund or carry out would adversely affect the delta smelt, the smelt population has collapsed since the state listing, de facto evidence that the CESA listing is inadequate to protect the species. No state recovery plan for the species exists. State listing has not prevented the delta smelt's decline and has not ensured that state projects such as the SWP protect the species. The CESA listing also does not protect delta smelt from the adverse effects of federal actions.

##### Federal Threatened Listing:

As demonstrated by the declining population trend since delta smelt was listed as threatened in 1993, a trend that has culminated in the recent catastrophic decline, as well as the extinction risk analysis discussed above, the protections currently afforded by the federal threatened listing, and implementation and enforcement of those protections, are clearly inadequate to protect delta smelt. During the past 13 years, many of the threats identified by USFWS in the original listing documents have not been eliminated or mitigated and have instead, in a number of instances, increased. Proposals to further increase the level of several key threats to the species (e.g., Delta water exports) have been approved by USFWS with "no jeopardy" biological opinions (USFWS 2004a, and see below). It is also apparent that the qualitative and quantitative recovery criteria specified for the species (Moyle et al. 1996) are insufficiently protective and inadequate to ensure continued existence of the species, particularly in the context of escalating threats to the species and inadequate regulatory protections.

##### 1995 SWRCB Water Quality Control Plan:

Overall water management operations in the Sacramento-San Joaquin Delta are regulated by the California State Water Resources Control Board, according to their 1995 Water

Quality Control Plan (WQCP) (SWRCB 1995). One subset of the WQCP's water quality objectives is specifically for the protection of fish and wildlife beneficial uses. Among other protections, these regulations limit Delta water exports in relation to Delta inflow (the Export/Inflow, or E/I, ratio), allowing up to 35% of total Delta inflow to be exported during the spring (February-June) and up to 65% to be exported during the rest of the year (July-January).<sup>9</sup> However, depending on inflow conditions, this can allow exports to be as high as (or higher than) 600,000 thousand acre-feet (AF) per month (or >10,000 cubic feet per second [cfs]), levels that, according to the analyses above, reliably correspond to low population levels for delta smelt.<sup>10</sup> For example, December-March exports have exceeded 2 million AF in nine of 39 years since 1967 (1983, 1988, 1989, 1990, 2000, 2002, 2003, 2004, and 2005). For those years, the delta smelt Recovery Index averaged 74, the level identified by the 2004 Biological Opinion to correspond to a "high level of concern" for the species and which is supposed to trigger additional protective actions (see Table 1). The Recovery Index fell below 74 in five of the nine years. By comparison, the Recovery Index exceeded 74 in 81% of years in which December-March exports were less than 1.5 million AF. Given that the Recovery Index has a threshold value of 239 that must be achieved in two of five sequential years for recovery of the species, this suggests that high export levels currently allowed by the WQCP and which have been implemented with increasing frequency during recent years (and which are predicted to occur even more frequently in the future; see 2004 Biological Opinion below) are incompatible with the recovery of delta smelt.

Export levels allowed under the WQCP also do not protect key areas of delta smelt critical habitat. Recent analyses using CDWR's Delta Simulation Model and its Particle Tracking Module show that most particles (which are thought to reasonably simulate larval and small juvenile delta smelt but be somewhat less accurate simulations for adult delta smelt) released into the southern and central Delta were lost within two weeks to entrainment at either the government water project diversions or the many unscreened agricultural diversions located in this area of the Delta (Figure 7). For particles released at Bacon Island in the southern Delta, virtually all the particles were entrained at E/I ratios >20%, a level well within the currently allowed operations. For particles released in the central Delta (Twitchell Island), the percentage entrained increased linearly with the E/I ratio. This shows that E/I ratios allowed by the current WQCP may be effectively eliminating large areas of the southern and central Delta as usable habitat for delta smelt spawning and early rearing (see Historic and Current Distribution above).


Although the E/I regulations established in 1995 with the WQCP prevented extreme water management operations such as those that occurred in the late 1980s and early 1990s, since that time (and since the delta smelt was listed under the ESA), seasonal E/I

---


<sup>9</sup> In the WQCP exports are defined as the combined exports of the federal Central Valley Project and the State Water Project. For a 31-day period during the spring (usually April 15-May 15), lower exports are required by the San Joaquin River Agreement as part of the Vernalis Adaptive Management Plan.

<sup>10</sup> Average Delta exports rates for the December-March period exceeded 10,000 cfs in 1990 and 2004. Since 2000, average December-March exports exceeded 9,000 cfs in four of the last six years (2000, 2002, 2003, and 2004).

ratios have increased by more than 100% for the winter period (from 14% in 1995-1999 to 30% in 2000-2004) and by more than 60% in the spring (from 16% to 26%) (Figure 8).



**Figure 7.** The relationship of the E/I ratio to the number of particles entrained (out of 5,000 particle released) over a series of Particle Tracking Model runs. Source: Herbold et al. 2005, Figure 8.



**Figure 8.** Trends in the Export/Inflow ratio (E/I) during the winter (December-March, upper panel) and spring (March-July, lower panel). The Water Quality Control Plan allows a maximum

E/I of 65% during the fall and winter (July-January) and 35% during the spring (February-June). Data source: California Department of Water Resources, Dayflow.

#### 2004 USFWS Biological Opinion:

A 2004 Formal and Early Endangered Species Consultation on the Coordinated Operations of the CVP and SWP and an Operations Criteria and Plan (USFWS 2004a) was intended to evaluate the impacts of the proposed future federal and state water management operations on delta smelt. The Consultation resulted in a June 2004 Biological Opinion by the U.S. Department of the Interior concluding that federal and state water exports from the Delta could increase without jeopardizing listed delta smelt.<sup>11</sup> The Consultation identified specific fish protection management tools, such as the CALFED Environmental Water Account (EWA), that will be used to mitigate the impacts of those actions on delta smelt using real-time management.<sup>12</sup> This regulatory mechanism provides inadequate protection for delta smelt for numerous reasons, including: 1) the effects of the proposed action are not adequately or correctly evaluated and adverse impacts are therefore seriously underestimated; and 2) the specified protective tools and management actions are known to be inadequate, not guaranteed to be available, and are not required to be implemented when requested by state and federal fish agencies for protection of the species.

The only evaluation of the impacts to delta smelt from the predicted increases in Delta water exports included in the 2004 Biological Opinion was based on estimates of increased incidental take. Increased take was calculated based on linear increases with increases in exports, an estimate that recent analyses by Herbold et al. (2005) suggest is seriously flawed (this analysis showed disproportionate increases in incidental take with recent increases in exports; see Figure 5). In addition, incidental take is known to be a poor predictor of either actual direct impacts or population level effects. The Biological Opinion lacks even basic statistical analyses of the effects of Delta exports on delta smelt population abundance. Given that statistically significant relationships between these important variables are detectable using the data that were available to the USFWS (and the BOR, which prepared the Biological Assessment) at the time the Biological Opinion was prepared (see Modification of Habitat, Population Level Effects above and Figure 6), this represents a failure to use best available science, a legal requirement for this type of regulatory mechanism. Further, the results of these types of analyses (as well as even a cursory review of recent abundance and export data) clearly show that export rates proposed in the described project are incompatible with stabilizing the population at levels above those identified in the Biological Opinion that trigger a “high level of concern”, much less recovering the species.<sup>13</sup>

---

<sup>11</sup> The June 2004 Biological Opinion was slightly revised and reissued in February of 2005.

<sup>12</sup> The Environmental Water Account (EWA) is a supply of water managed by the state and federal fish agencies to facilitate periodic reductions in SWP and/or CVP exports at times when ESA-listed fish species are near the facilities and vulnerable to entrainment. EWA water is used to compensate the SWP and CVP for reductions in deliveries that might result from the export curtailment.

<sup>13</sup> In the action evaluated by the Biological Opinion, average December-March exports are predicted to be >9,000-10,000 cfs in nearly all years.

The 2004 Biological Opinion uses a “Delta Smelt Risk Assessment Matrix” (DSRAM) to identify conditions under which delta smelt are at risk from water project operations (primarily entrainment loss at the Delta export facilities) and identifies “export reduction at one or both facilities” as one its protective “tools for change”.<sup>14</sup>

To implement protective actions recommended by the multi-agency Delta Smelt Working Group (DSWG, also specified and described in the Biological Opinion), the Biological Opinion relies on the CALFED Bay-Delta Program’s EWA to facilitate export reductions and to provide water resources to compensate for reduced deliveries that might result from the fish protection action. However, to date the EWA has been authorized and funded only on an annual basis and the likelihood of its continued existence beyond 2006 is unknown. In addition, the EWA’s utility for fish protection and the ability of the state and federal fish agencies to use it to implement specific protective actions as recommended by the multi-agency DSWG appears highly uncertain. In 2005, two out of three recommendations for protective actions made by the DSWG based on the risk criteria in the DSRAM were not implemented as recommended by the state and federal water project agencies (despite adequate water assets in the EWA) (Poage 2005). For example, in January of 2005 at least three of the risk criteria were exceeded and the monthly incidental take limit was nearly exceeded before an export reduction was implemented, an inexplicable delay given the high level of concern based on the then record low FMWT Index. When the reduction was finally implemented, neither the recommended export level nor duration was agreed to by the Water Operations Management Team (WOMT), who opted instead for a smaller and shorter export curtailment. Later in the spring, the DSWG-recommended export level for the 31-day Vernalis Adaptive Management Plan (VAMP) was rejected by the WOMT, who allowed for exports 50% greater than the recommended level. Three weeks later the SWP unilaterally discontinued its export curtailment before the VAMP period was concluded. The following summer, the abundance of juvenile delta smelt reached a record low and, when adult fish were surveyed in the fall, their numbers were also at new record lows.

#### Central Valley Project Improvement Act

The Central Valley Project Improvement Act (CVPIA), implemented in 1992, was intended to improve habitat conditions and reduce adverse direct and indirect impacts of CVP operations on fish and wildlife resources, with an emphasis on anadromous fish species. With the exception of small, short-duration water export reductions at the CVP export facility (usually timed to protect migrating juvenile salmonids), most of the habitat restoration and protective actions specified by the program (e.g., gravel restoration, stream flow enhancement, installation of fish screens and ladders) have been implemented outside the geographic range of the delta smelt and therefore provide neither benefit nor protection to the species.

---

<sup>14</sup> Page 98 in USFWS 2004a.



### CALFED Bay-Delta Program:

The CALFED Bay-Delta Program (CALFED) is a joint state-federal effort to restore the ecological health of the Bay-Delta system, protect and recover fish and wildlife species, increase water supply and reliability, and improve levee integrity and water quality. The delta smelt is identified by CALFED as a priority species. Within delta smelt habitat, CALFED has implemented three categories of restoration and protective actions intended, at least in part, to protect and recover the species: restoration of shallow water habitat; reduction of entrainment losses using the EWA; and installation of fish screens.

*Habitat restoration:* Restoration of shallow water habitat within the delta smelt's range has proceeded slowly and there is little evidence that the species derives benefit from the few small projects that have been completed (Brown 2003). There is no real evidence that access to shallow water habitat represents a limiting factor for the species (Bennett 2005).

*Export curtailments and the EWA:* The most recent Biological Opinion for delta smelt (USFWS 2004a) relies heavily on use of the EWA, a tool to facilitate periodic export curtailments and reduce entrainment losses that was developed by CALFED as a four-year experiment. The USFWS (2004b) cited the EWA and the 31-day-long spring VAMP export curtailment that is largely implemented using EWA water as examples of water management tools that are addressing specific needs of delta smelt. During the past five years, however, state and federal agencies have reduced by 300,000 to 400,000 acre-feet per year the amount of water required by the CALFED Record of Decision (ROD) to protect the Delta and fisheries under the EWA (Environmental Defense 2005). Moreover, the USFWS (2004b) also noted that the delta smelt population declined during the three years of EWA actions that were analyzed in their 5-year review. In 2005, after five years of EWA implementation, Delta exports during seasons when delta smelt are vulnerable to direct and indirect impacts have continued to increase and the abundance of juvenile and adult delta smelt population has dropped to critically low levels.

The EWA is now in its sixth year of implementation. The program has been subject to four consecutive formal reviews by an independent science panel and a fifth informal review late last year. Despite these exhaustive reviews, the state and federal fish agencies managing the EWA have presented no evidence to indicate that the EWA is an effective tool for mitigating the adverse impacts of Delta export operations or even for reducing entrainment loss of fish at the SWP and CVP facilities. Similarly, there is no evidence that the EWA provides population level benefits to any of the ESA-listed species that are the focus of its actions. The current EWA is clearly not providing adequate protection to delta smelt. The USFWS (2004b) reported that the EWA is currently being "resized" to account for proposed increases in water exports at the state and federal export facilities, however, the proposed future EWA actually has less water than the original version described in the CALFED ROD. Therefore, a "resized" EWA will not likely provide adequate protection for delta smelt either.

*Installation of fish screens:* The USFWS 5-year review (2004b) cited installation of two fish screens on water diversions within the range of delta smelt under the auspices of the CALFED Bay-Delta Program as representing “progress towards eliminating entrainment of delta smelt in unscreened diversions,” but acknowledged that over 1,800 unscreened or inadequately screened diversions still operate in delta smelt critical habitat. Plans to renovate and upgrade the fish louvers and screens at the SWP and CVP fish facilities, which entrain the vast majority of larval, juvenile and adult delta smelt, have been indefinitely postponed (SDFFF 2005)

#### Federal Listing of Other Species Within the Range of the Delta Smelt:

Three other fish species that periodically co-occur with delta smelt are listed under the ESA: winter-run Chinook salmon (*Oncorhynchus tshawytscha*), spring-run Chinook salmon (*O. tshawytscha*), and Central Valley steelhead (*O. mykiss*). A fourth species, green sturgeon (*Acipenser medirostris*), is proposed for listing. With the exception of periodic water export reductions using either the CVPIA or EWA (see above sections), most of the habitat restoration and protective actions for these species have been implemented outside the geographic range of the delta smelt and therefore provide neither benefit to nor protection of the species.

#### CEQA and NEPA:

The environmental review process under the California Environmental Quality Act, California Public Resources Code §21000 et. seq. (CEQA) should theoretically provide some protection to delta smelt. CEQA declares that it is the policy of the state to “(p)revent the elimination of fish or wildlife species due to man’s activities, ensure that fish and wildlife populations do not drop below self-perpetuating levels, and preserve for future generations representations of all plant and animal communities.” (California Public Resources Code, section 21001(c)). The CEQA process is triggered when discretionary activities of state agencies may have a significant effect on the environment. When the CEQA process is triggered, it requires full disclosure of the potential environmental impacts of proposed projects. The operative document for major projects is usually the Environmental Impact Report.

Theoretically, besides ensuring environmental protection through procedural and informational means, CEQA also has substantive mandates for environmental protection. The most important of these is the provision requiring public agencies to deny approval of a project with significant adverse effects when feasible alternatives or feasible mitigation measures can substantially lessen such effects. *Citizens for Quality Growth v. City of Mt. Shasta*, 198 Cal.App.3d 433, 440\_441 (1988); CA. Pub. Res. Code § 21002; 14 Cal. Code Regs. §§ 15002(a)(3), 15021(a)(2) and (c), 15041(c), 15364, 15370. In practice, however, this substantive mandate has not been implemented, especially with regard to protection of the delta smelt. In practice, alternatives that would protect the delta smelt and other wildlife are almost universally dismissed as “infeasible.” Mitigation, when required, is often ineffective or only marginally effective.

The National Environmental Policy Act (“NEPA”) also requires that federal agencies fully and publicly disclose the potential environmental impacts of proposed projects, but NEPA lacks even the minimal substantive provisions of CEQA.

## **5. Other Natural or Anthropogenic Factors**

### Reduction in Prey Food Organisms:

In recent years, untreated discharges of ship ballast water have introduced non-indigenous aquatic species to the Delta ecosystem (Carlton et al. 1990), several of which adversely affect the delta smelt and its habitat. The USFWS (2004b) noted that the discharge of any ballast water into San Francisco Bay is not prohibited and that compliance with a Coast Guard requirement to discharge ballast water before entering U.S. ports is voluntary. Without strictly enforced prohibitions on ballast water discharge in the Bay, additional introductions of non-native species can be expected to continue (Moyle et al. 1996, Moyle 2003). The USFWS (1993, 2004a, 2004b) discussed the impacts of an introduced clam (*Corbula amurensis*, formerly *Potamocorbula amurensis*) on the primary food of delta smelt and of non-native copepods that reduce food availability or feeding efficiency of delta smelt. The USFWS (1993, 2004a, 2004b) also discussed the altered plankton food web in the Delta that may have decreased the growth efficiency of delta smelt larvae.

### Toxic Substances:

The USFWS (1993, 2004a, 2004b) discussed the threat to delta smelt of poor water quality due to discharge and transport of agricultural and industrial chemicals from the Central Valley and Delta. Surveys of Delta waters have detected multiple pesticides and herbicides (Kuivila 2000; Houston et al. 2000). The USFWS (1993) noted that irrigation drain water can be harmful to other Delta fish larvae and embryos and to the planktonic food organisms for delta smelt. The SWRCB has designated all of the important water bodies in the delta smelt’s range as impaired by one or more contaminants, commonly including pesticides such as diazinon, chlorpyrifos, malathion, chlordane, DDT and dieldrin. There is growing evidence that fish species in the Delta are suffering direct mortality, and physiological and/or developmental impairment from the presence of toxic substances in the water and in the plankton upon which the smelt feed, and that their planktonic food supply may be depleted by periodic, highly concentrated pulses of herbicides and pesticides through the Delta (see USFWS 1996, 1999).

### Potential Loss of Genetic Integrity:

Species for which the population level falls below its effective population size are subject to inbreeding and genetic drift. This can result in depressed reproductive vigor and loss of genetic variation, reducing evolutionary fitness of the species and increasing its extinction risk. Using the effective population size criterion developed by Bennett (2003, 2005) and recognized by the USFWS (2004b), two consecutive years of a FMWT Index

less than 100, the delta smelt is now at risk for loss of its genetic integrity and at greater risk of extinction than when the species was listed under the ESA.

### **III. CONCLUSION**

The USFWS (2004b) recently concluded that the threats to delta smelt described in the original listing remain, including the destruction, modification, or curtailment of the delta smelt's habitat or range resulting from extreme freshwater outflow conditions (reduced outflow or high outflow), operations of the State and Federal water projects, and other water diversions. CDFG (2003) and Moyle (2002, 2003) concluded that these threats could result in the extinction of the delta smelt. In addition, any one of the many stochastic factors that affect delta smelt, such as predation, invasive species, change in food organisms, toxic substances, disease, competition, and entrainment losses to water diversions can cause their numbers to move towards extinction (Moyle 2002, 2003).


In 2003, the USFWS (2004b) concluded that new information concerning the delta smelt's population size and extinction probability indicated that the population was at risk of falling below an effective population size and therefore in danger of becoming extinct. Since the USFWS status review, the delta smelt population has indeed fallen below an effective population size and is therefore in danger of becoming extinct.

Recent population viability and extinction risk analyses indicate that, conservatively, the delta smelt could go extinct within the next 20 years and new analyses have demonstrated a statistically significant relationship between delta smelt population abundance and Delta export rates. Despite the availability of this new information regarding the likelihood of extinction of the species and the delta smelt's decline to record low levels for two consecutive years (2004 and 2005), neither the USFWS nor other agencies charged with protecting the delta smelt and its Delta habitat have taken major actions to reverse the decline. Indeed, state and federal agencies continue to proceed with plans to that would exacerbate current conditions for the delta smelt, including increasing export levels at the Delta pumping facilities and developing new water diversion and storage projects which will reduce Delta inflow and further degrade delta smelt habitat.

The warning signs could not be clearer: the delta smelt is at high risk of extinction. We urgently petition the USFWS to list the delta smelt as endangered, and to immediately implement measures to curtail threats to the species and improve environmental conditions in its Delta habitat.

#### IV. SIGNATURE PAGE

Submitted this 8th day of March, 2006



Jeff Miller  
Center for Biological Diversity  
1095 Market Street, Suite 511  
San Francisco, CA 94103  
Phone: (415) 436-9682 x303  
Fax: (415) 436-9683  
E-mail: [jmiller@biologicaldiversity.org](mailto:jmiller@biologicaldiversity.org)



Christina Swanson, Ph.D.  
The Bay Institute  
500 Palm Drive, Suite 200  
Novato, CA 94949  
Phone: (530) 756-9021  
Fax: (530) 756-9021  
E-mail: [swanson@bay.org](mailto:swanson@bay.org)

## V. BIBLIOGRAPHY OF LITERATURE CITED

Armor, C., R. Baxter, B. Bennett, R. Breuer, M. Chotkowski, P. Coulston, D. Denton, B. Herbold, K. Larsen, M. Nobriga, K. Rose, T. Sommer, and M. Stacey. 2005. Interagency Ecological Program synthesis of 2005 work to evaluate the Pelagic Organism Decline (POD) in the upper San Francisco Estuary. Available at: [http://www.science.calwater.ca.gov/workshop/workshop\\_pod.shtml](http://www.science.calwater.ca.gov/workshop/workshop_pod.shtml)

Bennett, W. A. 1995. Potential effects of exotic inland silversides on delta smelt. Interagency Ecological Program for the Sacramento-San Joaquin Estuary Newsletter. 8(1):4-6.

Bennett, W. A. 2003. The Population Ecology of Delta Smelt in the San Francisco Estuary. 44 pp.

Bennett, W. A. 2005. Critical assessment of the delta smelt population in the San Francisco Estuary, California. *San Francisco Estuary and Watershed Science* 3(2):1-71.

Brown, L. R. 2003. Will tidal wetland restoration enhance populations of native fishes? *San Francisco Estuary and Watershed Science*. Vol. 1, Article 2. 42 pp.

Bowen, M. D., B. B. Baskerville-Bridges, K. W. Frizell, L. Hess, C. A. Carp, S. M. Siegfried, and S. L. Wynn, 2004. Empirical and experimental analyses of secondary louver efficiency at the Tracy Fish Collection Facility, March 1996 to November 1997, Tracy Fish Facility Studies, Volume 11, U. S. Bureau of Reclamation, Mid Pacific Region, Denver Technical Service Center, 33 pp + Figures.

CALFED Bay-Delta Program. 2001. Delta Smelt and CALFED's Environmental Water Account: Summary of a Workshop held September 7, 2001 at the University of California, Davis. Sacramento, California.

California Department of Fish and Game (CDFG). 2003. Comment Letter on the Five Year Status Review of the Delta Smelt. 6 pp.

California Department of Water Resources (CDWR). 2003. Comment Letter on the Five Year Status Review of the Delta Smelt. 20 pp.

Carlton, J. T., J. K. Thompson, L. E. Schemel, and F. H. Nichols. 1990. Remarkable invasion of San Francisco Bay (California, USA) by the Asian clam *Potamocorbula amurensis*. I. Introduction and dispersal. *Mar. Ecol. Prog. Ser.* 66: 81-94.

Environmental Defense. 2005. Finding the water. New water supply opportunities to revive the San Francisco Bay-Delta ecosystem. Environmental Defense, Oakland, CA. 22 pp.

- Erkkila, L. F., J. F. Moffett, O. B. Cope, B. R. Smith, and R. S. Nelson. 1950. Sacramento-San Joaquin Delta fishery resources: effects of Tracy pumping plant and delta cross channel. U.S. Fish and Wildl. Serv. Spec. Sci. Rep. Fisheries 56. 109 pp.
- Fleming, K. and M. Nobriga. 2004. Status of delta smelt: multi-decadal trends in abundance, distribution, and water management. Proceedings of the 2004 CALFED Bay-Delta Program Science Conference. October 4-6, 2004. Sacramento, California.
- Herbold, B., C. Armor, R. Baxter, M. Chotkowski, P. Coulston, M. Nobriga, and T. Sommer. 2005. Historical patterns in salvage data. In: IEP synthesis of 2005 work to evaluate the pelagic organism decline (POD) in the upper San Francisco Estuary. Available at: [http://www.science.calwater.ca.gov/workshop/workshop\\_pod.shtml](http://www.science.calwater.ca.gov/workshop/workshop_pod.shtml)
- Herren, J. R. and S. S. Kawasaki. 2001. Inventory of water diversions in four geographic areas in California Central Valley. California Fish and Game Fish Bulletin 179:343-355.
- Houston, J. R, L. A. Allen, and K. M. Kuivila. 2000. Seasonal patterns and factors controlling the occurrence of dissolved pesticides in the Sacramento-San Joaquin Delta. Presented at CALFED Bay-Delta Program Science Conference, Oct. 3-5, 2000, Sacramento, CA. Abstract (#169), summary, and notes available at [www.iep.water.ca.gov/calfed/sciconf/2000/publications/](http://www.iep.water.ca.gov/calfed/sciconf/2000/publications/).
- Kimmerer, W. J. and J. J. Orsi. 1996. Changes in the zooplankton of the San Francisco Bay Estuary since introduction of the clam *Potamocorbula amurensis*. Pages 403-424. In J. T. Hollibaugh, ed. San Francisco Bay: the ecosystem. Pacific Division of the American Association for the Advancement of Science. San Francisco, California, USA.
- Kuivila, K. M. 2000. Pesticides in the Sacramento-San Joaquin Delta: State of Our Knowledge. Presented at CALFED Bay-Delta Program Science Conference, Oct. 3-5, 2000, Sacramento, CA. Abstract (#66), summary, and notes available at [www.iep.water.ca.gov/calfed/sciconf/2000/publications/](http://www.iep.water.ca.gov/calfed/sciconf/2000/publications/).
- Mager, R. 1996. Gametogenesis, reproduction, and artificial propagation of delta smelt, *Hypomesus transpacificus*. Ph.D. Dissertation. University of California, Davis.
- Matica, Z. and T. Sommer. 2005. Aquatic impacts of the Pittsburg and Contra Costa power plant. In: IEP synthesis of 2005 work to evaluate the pelagic organism decline (POD) in the upper San Francisco Estuary. Available at: [http://www.science.calwater.ca.gov/workshop/workshop\\_pod.shtml](http://www.science.calwater.ca.gov/workshop/workshop_pod.shtml)
- Miller, L. W. 2000. The tow-net survey for delta smelt revisited. Interagency Ecological Program for the Sacramento-San Joaquin Estuary Newsletter. 13(1):37-44.

Moon, G. E, K. M. Kuivila, and J. L. Orlando. 2000. Exposure of Delta Smelt to Dissolved Pesticides During Larval and Juvenile Stages in 1998 and 1999. Abstract #183 presented at CALFED Bay-Delta Program Science Conference, Oct. 3-5, 2000, Sacramento, CA. Summary and notes available at [www.iep.water.ca.gov/calfed/sciconf/2000/publications/](http://www.iep.water.ca.gov/calfed/sciconf/2000/publications/).

Moyle, P. B. 2003. Comment Letter on the Five Year Status Review of the Delta Smelt. 4 pp.

Moyle, P. B. 2002. Inland Fishes of California. University of California Press, Berkeley, California, USA.

Moyle, P. B. and B. Herbold. 1989. The delta smelt, *Hypomesus transpacificus*. Unpublished final report prepared for U.S. Fish and Wildlife Service, Sacramento Field Office, Habitat Conservation Division, Sacramento, California. 42 pp.

Moyle, P. B., B. Herbold, D. E. Stevens, and L. W. Miller. 1992. Life history and status of delta smelt in the Sacramento-San Joaquin estuary, California. Transactions of the American Fisheries Society 121:67-77.

Moyle, P. B., R. Pine, L. R. Brown, C. H. Hanson, B. Herbold, K. M. Lentz, L. Meng, J. J. Smith, D. A. Sweetnam, and L. Winternitz. 1996. Sacramento-San Joaquin Delta native fishes recovery plan. Portland (OR): U.S. Fish and Wildlife Service.

National Marine Fisheries Service (NMFS) and U.S. Fish and Wildlife Service (USFWS). 1974. Memorandum of Understanding between the U. S. Fish and Wildlife Service, United States Department of the Interior and the National Marine Fisheries Service, National Oceanic and Atmospheric Administration, United States Department of Commerce regarding jurisdictional responsibilities and listing procedures under the Endangered Species Act of 1973.

Nobriga, M, Z. Matica, and Z. Hymanson (2002) Assessing fish entrainment vulnerability to agricultural irrigation diversions: a comparison among native and non-native species. Interagency Ecological Program for the Sacramento-San Joaquin Estuary Newsletter. 15(3):35-44.

Poage, V. 2005. Environmental Water Account expenditures for the protection of delta smelt in Water Year 2005. Presented at the 2005 Annual Review of the Environmental Water Account, Dec. 7-8, 2005, Sacramento, CA, and available at: <http://science.calwater.ca.gov/workshop/ewa.shtml>.

Radtke, L. D. 1966. Distribution of smelt, juvenile sturgeon and starry flounder in the Sacramento-San Joaquin Delta. Fish Bulletin 136. Sacramento (CA):California Department of the Fish and Game. P. 115-129.



San Luis & Delta-Mendota Water Authority. 2002. The Delta Smelt and the State of the Science. 5 pp + Appendices.

Simi, J. and C. Ruhl. 2005. Summary of Delta hydrology data water years 1985-2004. In: IEP synthesis of 2005 work to evaluate the pelagic organism decline (POD) in the upper San Francisco Estuary. Available at:  
[http://www.science.calwater.ca.gov/workshop/workshop\\_pod.shtml](http://www.science.calwater.ca.gov/workshop/workshop_pod.shtml)

South Delta Fish Facilities Forum (SDFFF). 2005. South Delta Fish Facilities Forum Co-Chairs Report: Some Policy Conclusions, April 20, 2005. Available at:  
<http://calwater.ca.gov/Programs/Conveyance/SDFF/SouthDeltaFishFacilitiesForum.shtml>.

Souza, K. K. Hieb, K. Fleming, M. Bryant, and R. Baxter. 2005. Apparent growth rates of pelagic fishes and relation to abundance. In: IEP synthesis of 2005 work to evaluate the pelagic organism decline (POD) in the upper San Francisco Estuary. Available at:  
[http://www.science.calwater.ca.gov/workshop/workshop\\_pod.shtml](http://www.science.calwater.ca.gov/workshop/workshop_pod.shtml)

Stanley, S. E., P. B. Moyle and H. B. Shaffer. 1995. Allozyme analysis of delta smelt, *Hypomesus transpacificus* and longfin smelt, *Spirinchus thaleichthys* in the Sacramento-San Joaquin Estuary, California. *Copeia* 1995:390-396.

State Water Resources Control Board (SWRCB). 1995. Water Quality Control Plan for the San Francisco Bay/Sacramento-San Joaquin Delta estuary. State Water Resources Control Board, California Environmental Protection Agency. 95-1WR, May 1995.

Stevens, D. E. and L. W. Miller. 1983. Effects of river flow on abundance of young chinook salmon, American shad, longfin smelt, and Delta smelt in the Sacramento-San Joaquin river system. *N. Am. J. Fish. Manage.* 3:425-437.

Swanson, C. 2005. Effects of exports on delta smelt population abundance – preliminary analyses. Presented at the 2005 Annual Review of the Environmental Water Account, Dec. 7-8, 2005, Sacramento, CA, and available at:  
<http://science.calwater.ca.gov/workshop/ewa.shtml>.

Swanson, C., T. Reid, P. S. Young, and J. J. Cech, Jr. 2000. Comparative environmental tolerances of threatened delta smelt (*Hypomesus transpacificus*) and introduced wakasagi (*H. nipponensis*) in an altered California estuary. *Oecologia* 123:384-390.

Swanson, C., P. S. Young, and J. J. Cech, Jr. 2005. Close encounters with a fish screen: integrating physiological and behavioral results to protect endangered species in exploited ecosystems. *Transactions of the American Fisheries Society* 134:1111-1123.

Teh, S. J. 2005. Histopathological evaluation of starvation and/or toxic effects on pelagic fishes. In: IEP synthesis of 2005 work to evaluate the pelagic organism decline (POD) in the upper San Francisco Estuary. Available at:  
[http://www.science.calwater.ca.gov/workshop/workshop\\_pod.shtml](http://www.science.calwater.ca.gov/workshop/workshop_pod.shtml)

Trenham, P. C., H. B. Shaffer, and P. B. Moyle. 1998. Biochemical identification and assessment of population subdivision in morphologically similar native and invading smelt species (*Hypomesus*) in the Sacramento-San Joaquin estuary, California. *Transactions of the American Fisheries Society* 127:417-424.

U.S. Fish and Wildlife Service (USFWS). 1993. Determination of Threatened Status for the Delta Smelt. 58 FR 12854 12864, March 5, 1993.

U.S. Fish and Wildlife Service (USFWS). 1994. Critical Habitat Determination for the Delta Smelt. 59 FR 65256 65279, December 19, 1994.

U.S. Fish and Wildlife Service (USFWS). 1996. Recovery Plan for the Sacramento-San Joaquin Delta Native Fishes.

U.S. Fish and Wildlife Service (USFWS). 1999. Determination of Threatened Status for the Sacramento Splittail. 64 FR 5963, 5974-80, Feb. 8, 1999.

U.S. Fish and Wildlife Service (USFWS). 2004a. Formal and Early Section 7 Endangered Species Consultation on the Coordinated Operations of the Central Valley Project and State Water Project and Operational Criteria and Plan. U. S. Fish and Wildlife Service, Sacramento Fish and Wildlife Office, Sacramento, California.

U.S. Fish and Wildlife Service (USFWS). 2004b. 5-Year Review, *Hypomesus transpacificus* (delta smelt). U. S. Fish and Wildlife Service, Sacramento Fish and Wildlife Office, Sacramento, California.

U.S. Geological Survey (USGS). 2003. Editor's Comments-Peer Review of the 2002 San Luis and Delta-Mendota Water Authority White Paper "The Delta Smelt and the State of the Science."